

Code: 30217

THIODAN KOLODUST[®] 3-25 DUST

50 Pounds Net Weight

DIRECTIONS FOR USE

Apply when insects or disease first appear and repeat as necessary. For insect or disease control beyond the use limitations of this product use a suitable material. May be applied with ground or airplane application equipment.

APPLES to control: Apple Aphid use 50-80 pounds per acre. Do not apply within 21 days of harvest nor apply more than 3 applications during the fruiting period. With no limitations as to the number of applications do not apply within 30 days of harvest. Do not feed pomace from treated apples to livestock. Sulphur may injure certain varieties, check tolerance to sulphur prior to use.

GRAPES to control: Powdery Mildew, Grape Leafhopper, Grape Leafroller (Calif. Only) use 32-50 pounds per acre. For Leafroller control repeat applications may be necessary. Do not use on Concord variety of grapes as severe injury is likely to occur. Do not apply within 7 days of harvest. Apply with ground application equipment only.

PEACHES AND NECTARINES to control: Green Peach Aphid, Peach Twig Borer (Calif. Only) and Peach Silvermite (Calif. Only) use 50-80 pounds per acre. Do not make more than 2 applications during the fruiting season nor apply within 30 days of harvest.

PLUMS AND PRUNES to control: Peach Twig Borer (Calif. Only) use 50-80 pounds per acre. Do not apply within 7 days of harvest.

FILBERTS to control: Filbert Aphid, Filbert Leafroller use 50-65 pounds per acre. May be applied up to the day of harvest.

BROCCOLI, CABBAGE to control: Cabbage Aphids, Cabbage Looper, Diamond-back Moth Larvae, Flea Beetle, Harlequin Bug, Stink Bug, Imported Cabbage Worm use 25-35 pounds per acre. Do not apply within 7 days of harvest.

COTTON to control: Cotton Stink Bug, Lygus Bug, Bollworm, Cabbage Looper, Thrips, Cotton Leaf Perforator use 32-50 pounds per acre. Do not apply after bolls begin to open. Do not graze meat or dairy animals in treated fields.

HEAD LETTUCE to control: Cabbage Looper, Diamond-back Moth Larvae, Green Peach Aphid, Imported Cabbage Worm use 32-50 pounds per acre. Do not apply more than 3 applications after treatment. Do not apply within 14 days of harvest. Remove outer leaves at harvest. Do not crop lettuce for livestock.

PEPPERS to control: Flea Beetle, Green Peach Aphid, Pepper Maggot, Hornworm use 32-50 pounds per acre. At the 35 pound rate do not apply within 1 day of harvest. At a rate above 35 pounds do not apply within 4 days of harvest.

POTATOES to control: Aphid, Leafhopper, Flea Beetle, Colorado Potato Beetle, Potato Tolerworm, Green Stinkbug, Southern Armyworm, Flea Psyllid, False Chinch Bug use 32-50 pounds per acre. Do not apply within 14 days of harvest.

SAFFLOWERS to control: Green Peach Aphid use 32-50 pounds per acre. Do not apply after flower heads open.

SEED PEAS to control: Pea Aphid and Pea Weevil use 25-35 pounds per acre. Do not feed treated vines or other parts to meat or dairy animals.

FUNGICIDE - INSECTICIDE
U.S.D.A. REGISTRATION NO. 279-1710

ACTIVE INGREDIENTS:	BY WT.
Endosulfan hexachlorocyclopentadiene	3.00%
2,4,3-benzodioxathiepin oxide	21.00%
Sulphur	76.00%
INERT INGREDIENTS:	100.00%

Thiodan is a registered trademark of Canadian Hoechst Ltd.-U.S. Patent No. 2,799,685

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water, for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and feedstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or domestic animals. Workers entering treated areas on the day of application should wear protective clothing.

USAGE CAUTION

This product is poisonous to fish and wildlife. Keep out of ponds, streams, and other aquatic areas. Do not apply in a manner not specified on the label.

Dispose of Wastes by burying in non-crop lands away from water supplies. Bag should be disposed of by burying with wastes or by burning. Keep Out of Smoke!

Application of water after treatment will not reduce the toxicity of the dust. Do not apply to crops or plants which are under stress or which are susceptible to frost damage.

Do not apply to water or to areas where it may drift into water. Do not apply to areas where it may drift into water. Do not apply to areas where it may drift into water.

STRAWBERRIES to control: Powdery Mildew, Meadow Spittlebug and Strawberry Aphid use 32-50 pounds per acre. Do not apply within 4 days of picking. For multiple applications during the fruiting period, do not apply at intervals of less than 15 days or more than twice during any 35 day period.

SUGARBETTES to control: Green Peach Aphid, Sugar Beet Weevil and Meadow Spittlebug use 32-50 pounds per acre. Do not feed treated tops to livestock.

TOMATOES to control: Aphid, Flea Beetle, hornworm, Colorado Potato Beetle, Tomato Psyllid, whitefly, Stinkbug, Tomato Fruitworm, Cabbage Looper, Yellow Striped Armyworm and Tomato Russet Mite use 32-50 pounds per acre. Do not apply within 1 day of harvest.

ORNAMENTALS (Greenhouse and Out-of-door) to control: Powdery Mildew, Aphid and Whitefly. Dust lightly but uniformly on both upper and lower leaf surfaces.

DRIED BEANS to control: Black Bean Aphid, Bean Leaf Skeletonizer and Mexican Bean Beetle use 32-50 pounds per acre. Do not apply within 40 days of harvest. Do not feed treated screenings to livestock. Sulphur may injure certain bean varieties, check tolerance to sulphur prior to use.

SUCCULENT BEANS to control: Black Bean Aphid, Bean Leaf Skeletonizer, Cucumber Beetle, Green Stinkbug, Mexican Bean Beetle use 32-50 pounds per acre. May be applied up to 3 days before harvest. Do not apply within 3 days of harvest. Use only on beans which are to be harvested by cutting. Do not feed treated screenings to livestock. Sulphur may injure certain bean varieties, check tolerance to sulphur prior to use.

BRUSSELS SPROUTS AND CABBAGE to control: Cabbage Aphid, Cabbage Looper, Imported Cabbage Worm, Diamond-back Moth Larvae, Flea Beetle, Harlequin Bug and Imported Cabbage Worm use 25-35 pounds per acre. Do not apply within 14 days of harvest.

USAGE CAUTION: Do not allow this material to drift onto neighboring crop or non-crop areas or into a stream or pond at any time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

Dealers Should Sell in Original Packages Only.


NOTICE: Niagara warrants, to the extent of the purchase price, that the goods will conform to the ingredients statement set forth on the label. EXCEPT AS SO WARRANTED, THE GOODS ARE SOLD "AS IS." NIAGARA MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED COVERING THE GOODS. Buyer or user assumes all risk and liability arising out of or connected with possession, handling, storage or use of the goods and accepts and uses the goods on these conditions. Niagara shall not be held responsible in any manner for personal injury, property damage or other loss of any kind including incidental or consequential damages, relating to possession, handling, storage or use of the goods, whether or not in accordance with directions, regardless of claimed negligence or fault of Niagara. No person is authorized to vary or waive anything contained in this notice.

Made in U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



279-1710



Dealers Should Sell in Original Packages Only.

USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara: (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 30 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK



FRESNO, CALIF. — RICHMOND, CALIF.
RIVERSIDE, CALIF. — YAKIMA, WASHINGTON