

Code:

50 Pounds Net Weight

SUNLAND WEEDABLES

ON APPLE...
Apply when disease threatens during rain periods and repeat at 7-10 day intervals.

On ALMONDS for Brown Rot and Scab control use 6 pounds per 100 gallons of water in pre-bloom, bloom and post-bloom periods.

On GRAPES for Powdery Mildew control use 2-5 pounds per 100 gallons of dilute spray. Use as a spray beginning when shoots are 6-8 inches long and repeat at 7 to 10 day intervals.

On ALMONDS for Brown Rot and Scab control use 6 pounds per 100 gallons of water in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7-10 day intervals.


On PLUMS, PRUNES, PEACHES, NECTARINES and CHERRIES for control of Powdery Mildew, Cherry Leaf Spot, Prune Rust and Brown Rot use at the

ACTIVE INGREDIENT: Sulphur By Wt. 95.00%
INERT INGREDIENTS 5.00%
100.00%

The material in this bag is guaranteed to pass 95% through a 325 mesh screen.

CAUTION

KEEP OUT OF REACH OF CHILDREN



Manufactured in U.S.A. by
SUNLAND WEEDABLES
RICHMOND, CALIF.
YACON, WASH.
New York: Mid-West, N.Y.

rate of 6 pounds per 100 gallons of water...
in the pre-bloom, bloom and post-bloom periods.
Apply when disease threatens during rain periods and repeat at 7-10 day intervals.

On STRAWBERRIES and BUSH BERRIES for control of Powdery Mildew use 2 pounds per 100 gallons of water. Apply when disease first appears and repeat as necessary.

On CITRUS for control of Citrus Flat Mite, Tuma Spider Mite, Six-spotted Mite and Citrus Rust Mites use 4-6 pounds per 100 gallons of water (a minimum of 60 pounds per acre). Apply when mites appear and repeat as necessary.

MIXING DIRECTIONS - Add the material to water until the agitator starts to run. With the agitator running add the required amount of Sunland Weedable Sulphur. Stop filling and keep agitator running.

USAGE CAUTION

Applications of materials containing sulphur should not be made prior to or during periods of excessively high summer temperatures or within 2 weeks of an oil spray. On citrus do not make an oil application within 60 days prior to or after a sulphur application.

Do not use on or allow this material to drift onto melons, squash, cucumbers, apricots or other plants which will not tolerate sulphur.

Cameraman's Note
Poor Copy