

**CODE: 30749**

**50 POUNDS NET WEIGHT**

# DUSTING SULPHUR

## DIRECTIONS FOR USE

On **APPLES, PEARS** for control of Scab, on **ALMONDS, CHERRIES, PEACHES, PLUMS AND PRUNES** for the control of Brown Rot, Cherry Leaf Spot and Prune Rust and on **APPLES, NECTARINES AND PEACHES** for the control of Powdery Mildew use at the rate of 40 to 50 pounds per acre in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. On Pears do not apply after the fruit begins to turn down. Sulphur may injure certain varieties of apples, check tolerance prior to using.

On **GRAPES** for control of Powdery Mildew use 8 to 15 pounds per acre. Begin applications when shoots are 6 to 8 inches long and repeat as necessary. On **COTTON**: For control of Strawberry Mite (Atlantic Mite) apply 35 to 40 lbs. per acre when mites first appear and repeat as necessary.

On **STRAWBERRIES AND BUSHBERRIES**: For control of Powdery Mildew use 10 to 20 pounds per acre. Apply when disease first appears and repeat as necessary

On **CITRUS**: For control of Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Six Spotted Mite and Citrus Rust Mite, use at the rate of 100 pounds per acre. Apply when pests first appear and repeat as necessary.

**VEGETABLE AND FIELD CROPS**: Apply recommended rate when pests or disease first appear and repeat as necessary. For disease control keep new growth protected.

## PESTICIDE

U.S.D.A. REGISTRATION NO. 279-1721

<b>ACTIVE INGREDIENT:</b>	<b>By Wt.</b>
Sulphur .....	98.00%
<b>INERT INGREDIENTS .....</b>	<b>2.00%</b>
	<b>100.00%</b>

The material in this bag is guaranteed to pass 95% through a 325 mesh screen.

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**

On **BEANS**: Powdery Mildew, Rust, Thrips, Sp. Mites—Apply 15 to 30 pounds per acre. For control start applications as soon as the first leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties. Check tolerance prior to application.

On **CARROTS**: Petrobia Mite — Apply 30 to 40 pounds per acre.

On **ONIONS**: Petrobia Mite — Apply 20 to 30 pounds per acre.

On **PEAS**: Powdery Mildew — Apply 15 to 20 pounds per acre.

On **TOMATOES**: Tomato Russet Mite — Apply 20 to 40 pounds per acre.

On **TURNIPS AND RUTABAGAS**: Powdery Mildew — Apply 15 to 20 pounds per acre.

On **ORNAMENTALS**: Rust, Powdery Mildew—Dust for complete coverage.

## CAUTION

Application of dusting and spraying materials containing sulphur should not be made just prior to or during periods of excessively high temperatures. Do not make applications within two weeks of an oil spray.

On Citrus do not make an oil application within 60 days prior to or after a sulphur application.

Do not use on or allow this material to drift on melons, squash, cucumbers, apricots or other plants which will not tolerate sulphur.

50 POUNDS NET WEIGHT

# DUSTING SULPHUR

### FOR USE

control of Scab, on  
PEACHES, PLUMS AND  
Brown Rot, Cherry Leaf  
APPLES, NECTARINES  
control of Powdery Mildew  
10 pounds per acre in  
pre-bloom periods. Apply  
during rain periods and  
dormant periods. On Pears do not  
turn down. Sulphur  
of apples, check toler-

powdery Mildew use 8 to  
10 applications when shoots  
appear repeat as necessary.  
control of Strawberry Mite (At-  
tacks) per acre when mites  
appear necessary.

**RASPBERRIES;** For con-  
trol 10 to 20 pounds per  
acre at first appears and repeat

control of Citrus Thrips, Citrus Flat  
leaf Spotted Mite and Cit-  
rus Mite rate of 100 pounds per  
acre at first appear and repeat as

**CROPS:** Apply recom-  
mended rate first appear  
of disease control keep

### PESTICIDE

U.S.D.A. REGISTRATION NO. 279-1721

ACTIVE INGREDIENT:	By Wt.
Sulphur.....	98.00%
INERT INGREDIENTS.....	2.00%
	<hr/> 100.00%

The material in this bag is guaranteed to pass 95% through a 325 mesh screen.

### CAUTION

KEEP OUT OF REACH OF CHILDREN

106

On BEANS: Powdery Mildew, Rust, Thrips, Spider Mites—Apply 15 to 30 pounds per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties. Check tolerance prior to application.

On CARROTS: Petrobia Mite — Apply 30 to 35 pounds per acre.

On ONIONS: Petrobia Mite — Apply 20 to 30 pounds per acre.

On PEAS: Powdery Mildew — Apply 15 to 20 pounds per acre.

On TOMATOES: Tomato Russet Mite — Apply 30 to 40 pounds per acre.

On TURNIPS AND RUTABAGAS: Powdery Mildew — Apply 15 to 20 pounds per acre.

On ORNAMENTALS: Rust, Powdery Mildew—Dust for complete coverage.

### CAUTION

Application of dusting and spraying materials containing sulphur should not be made just prior to or during periods of excessively high temperatures. Do not make applications within two weeks of an oil spray.

On Citrus do not make an oil application within 60 days prior to or after a sulphur application.

Do not use on or allow this material to drift onto melons, squash, cucumbers, apricots or other plants which will not tolerate sulphur.

279-1721

On STRAWBERRIES: For control of Powdery Mildew use 10 to 20 pounds per acre. Apply when disease first appears and repeat as necessary.

On CUCUMBERS: For control of Cucurbit Powdery Mildew, Yuma Spider Mite, Silver Mite, and Cucumber Rust Mite use at the rate of 10 pounds per acre. Apply when pests first appear and repeat as necessary.

**VEGETABLE AND FIELD CROPS:** Apply recommended rate when pests or disease first appear and repeat as necessary. For disease control keep new growth protected:

On BEANS: Powdery Mildew, Rust, Thrips, Spider Mite—Apply 15 to 30 pounds per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties. Check tolerance prior to application.

On CARROTS: Petrobia Mite—Apply 30 to 50 pounds per acre.

On ONIONS: Petrobia Mite—Apply 10 to 15 pounds per acre.

On PEAS: Powdery Mildew—Apply 10 to 15 pounds per acre.

On TOMATOES: Tomato Rust Mite—Apply 20 to 40 pounds per acre.

On TURNIPS AND RUTABAGAS: Powdery Mildew—Apply 15 to 20 pounds per acre.

On ORNAMENTALS: Rust, Powdery Mildew—Dust for complete coverage.

**ALFALFA:** To control Spider Mites use 10 to 20 pounds per acre. Begin applications when mites first appear. May be applied by ground or air. Use no nitrogen fertilizer by air.

**ASPARAGUS:** To control Spider Mites use 10 to 20 pounds per acre. Apply when mites first appear and repeat as necessary.

**ON MELONS:** For Mildew control apply 25 to 50 pounds per acre by air or ground equipment. Do not apply this material to sulphur sensitive varieties.

by air.

**ASPARAGUS:** To control Spider Mites use 30 to 40 pounds per acre. Apply when mites first appear and repeat as necessary.

**ON MELONS:** For Mildew control apply 25 to 50 pounds per acre by air or ground equipment. Do not apply this material to sulphur sensitive varieties.

**ON CUCUMBERS:** To control Powdery Mildew apply 15 to 20 pounds per acre by air or ground. Repeat applications in 7 to 10 days if necessary.

**SUGAR BEETS:** To control Spider Mites use 30 to 40 pounds per acre. Begin treatments when mites become damaging. May be applied by ground or air. Use the higher rate by air. Use in hot weather may cause burn but injury is less severe if sulphur is applied following an irrigation.

**USAGE CAUTIONS**

Application of dusting and spraying material containing sulphur should not be made just prior to or during periods of excessively high temperatures. Do not make applications within two weeks of an oil spray.

On Citrus do not make an oil application within 60 days prior to or after a sulphur application.

Do not allow this material to drift onto sulphur sensitive crops.

**DISPOSE OF WASTE:** by burying in non-crop land away from water supplies. Empty containers should be buried with wastes or burned. Keep out of reach.



**MANUFACTURED IN U.S.A. BY**

**NIAGARA CHEMICAL DIVISION  
FMC CORPORATION**

WINDSOR, N. Y.  
PERDUE, CALIFORNIA  
RICHMOND, CALIF.    YACOMA, WASHINGTON  
MADE IN U.S.A.

**Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.**

### **DEALER SHOULD SELL IN ORIGINAL PACKAGES ONLY**

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:

**WARRANTY:** Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

**DAMAGES:** Buyer's or user's exclusive remedy for damages for breach of warranty, or negligence attributable to Niagara, is the return of the product. Damages shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 90 days of discovery by buyer or user, failing which Niagara or user waives any claim for such damage.

**EXCUSE FROM NON-CONTROLLABLE CONDITIONS:** Because of certain critical conditions created or increased by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Cameraman's Note  
Poor Copy