

107
1201

ACCEPTED
JAN 5 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 279-1721

Code 30749

Lbs. Net Wt.

Dusting Sulfur

Fungicide-Insecticide

EPA Reg. No. 279-1721 ZB

EPA Est., 279-

Active Ingredient:	By Wt.
Sulfur.....	98.00%
Inert Ingredients:.....	2.00%
	100.00%

The material in this bag is guaranteed to pass 95% through a 325 mesh screen.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from heat, sparks, or flame. Do not smoke while applying this product.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution *inhaled or absorbed through skin.*
Harmful if swallowed. *Avoid breathing dust or spray mist.*
Avoid contact with eyes, skin and clothing.
Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

If **swallowed**: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
If in **eyes**: Flush with plenty of water. Get medical attention if irritation persists.
If on **skin**: Wash with plenty of soap and water. *Get medical attention.*
For Emergency Assistance Call 716-735-3765
See other panels for additional precautionary information.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.
Do not enter treated areas for 24 hours unless appropriate protective clothing is worn.
Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.
Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.
Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION

Area treated with Dusting Sulfur on (insert date). Do not enter treated areas for 24 hours unless appropriate protective clothing is worn (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

7130 R

721
Page 1 of 3

(see attachment)

STORAGE AND DISPOSAL

Pesticide Storage

Keep away from fire and sparks

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Application of dusting and spraying materials containing sulfur should not be made just prior to or during periods of excessively high temperatures. Do not make applications within two weeks of an oil spray.

On Citrus do not make an oil application within 60 days prior to or after a sulfur application.

Apples, Pears: Scab, Almonds, Cherries, Peaches, Nectarines, Plums, Prunes: Brown Rot, Cherry Leaf Spot (Cherries only), Peach Silver Mite and Rust. **Apples, Nectarines, Peaches:** Powdery Mildew—Use 40 to 50 pounds per acre in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. On Pears do not apply after the fruit begins to turn down. Sulfur may injure certain varieties of apples, check tolerance prior to using

Citrus: Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite and Citrus Rust Mite—Use 100 pounds per acre. Apply when pests first appear and repeat as necessary.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and or storage of this material when such use and/or handling and or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

279-1721
Page 2 of 3

FMC—FMC Trademark

→ Pesticides: Citrus Flat Mite - Use 50 pounds per acre. Apply when mites first appear and repeat as necessary. May be applied by ground or air.

3

Dusting Sulfur Fungicide-Insecticide
Code 30749

Grapes: To control Powdery Mildew -- Use 8 to 15 lbs. of Dusting Sulfur per acre. Begin applications when shoots are 6 to 8 inches long and repeat at 7 to 10 day intervals. Concord and other Labrusca-type grapes may be injured by sulfur.

Blackberries, Boysenberries, Gooseberries, Loganberries, Raspberries, Strawberries: To control Powdery Mildew -- Use 10 to 20 lbs. of Dusting Sulfur per acre. Apply when disease first appears and repeat at 7 to 10 day intervals.

Vegetable, Field and Ornamental Crops: Apply recommended rates of Dusting Sulfur when pests or disease first appear and repeat as necessary. For disease control keep new growth protected.

Beans: To control Powdery Mildew, Rust, Thrips and Spider Mites -- Use Dusting Sulfur at 15 to 30 pounds per acre. For Rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties. Check tolerance prior to application.

Carrots: To control Petrobia Mite -- Use Dusting Sulfur at 30 to 35 pounds per acre.

Cotton: To control Strawberry Mite -- Use Dusting Sulfur at 35 to 40 pounds per acre. Apply when mites first appear and repeat as necessary.

Onion: To control Petrobia Mite -- Use Dusting Sulfur at 26 to 30 pounds per acre.

Ornamentals (flowers, shrubs, trees): To control Rust, and Powdery Mildew -- Use Dusting Sulfur at a rate that will completely cover foliage and repeat at 5 - 10 day intervals.

Peas, Turnips, Rutabagas: To control Powdery Mildew -- Use Dusting Sulfur at 15 to 20 lbs. per acre. Repeat at 7 to 14 intervals.

Tomatoes: To control Tomato Russet Mite -- Use Dusting Sulfur at 30 to 40 pounds per acre.

279-1721
Page 3 of 3