

Code 453

Gallons Net Contents

Ethion Superior 70 Oil

Insecticide

EPA Reg No 279-1731

EPA Est. 279-

Active Ingredients	By Wt
Ethion	2.0%
Petroleum Oil	94.0%
Inert Ingredients	4.0%
	100.0


100000 Tetraethylthiopyrophosphate
 Contains 0.15 lbs. Ethion per gallon
 Petroleum Oil Specifications
 Unsulfonated Residue (Min)
 0.50%

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

For Emergency Assistance Call 716-735-3765
 See Other Panels for Additional Precautionary Information

FMC

FMC Corporation
 Agricultural Chemical Group
 2000 Market Street
 Philadelphia Pennsylvania 19103



PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution
 Harmful if swallowed. May be absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not store below 32°F. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not reuse this container. To dispose of container, triple-rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. Do not cut or weld this container. Consult Federal, State or local disposal authorities for approved alternative procedures.

Spray in warm days in late winter or early spring and before the buds separate into the clusters. Do not spray in freezing weather or early spring freezes after the buds have burst to any extent. Do not spray when buds have separated in the clusters.

Do not apply within 2 weeks of a later spray or within 7 days of Captan.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if when this product is added to the water in the spray tank a uniform milk white spray mixture results provided the spray mixture is being agitated. If the spray mixture after adding the oil to the spray tank with the agitator running is not milk white, do not use the resulting spray or the remainder of this product.

Fill the spray tank about one third full of water. Start the agitator and pour in the required amount of Ethion Superior 70 Oil. If other spray materials are to be used with the oil, mix the other materials thoroughly with the water in the spray tank first and then add the Ethion Superior 70 Oil as the filling of the tank is being completed. Do not fill in undisturbed form.

When the agitator has been started or the tank stopped for any length of time, spray in the ground for a few moments until the spray appears creamy white. This is desirable also where there are drips or leaks in the pipelines of stationary nozzles. All the material in stationary nozzles should be sprayed at each day.



Almonds, Peaches: Clover (Brown Almond) Mite, European Red Mite. To control overwintering Mite eggs use 2 gallons per 100 gallons of water. Oystershell Scale, San Jose Scale. Use 2 gallons per 100 gallons of water. Paratonia (Olive) Scale. Use 3 gallons per 100 gallons of water. Apply during dormant or delayed dormant period. Do not graze livestock in treated groves.

Apples: Apple Aphid, European Red Mite, Oystershell Scale, Rosy Apple Aphid, San Jose Scale. Use 2 gallons per 100 gallons of water. Make a single application during the delayed dormant period. Do not apply later than the delayed dormant period.

Apricots: Oystershell Scale. Use 2 gallons per 100 gallons of water. Apply during dormant or delayed dormant period.

Cherries (Sweet & Tart): Black Cherry Aphid, European Red Mite, Forbes Scale, Putnam Scale, San Jose Scale. Use 2 gallons per 100 gallons of water. Apply during delayed dormant period.

Nectarines: Clover (Brown Almond) Mite, European Red Mite. To control overwintering Mite eggs use 2 gallons per 100 gallons of water. San Jose Scale. Use 2 gallons per 100 gallons of water. Paratonia (Olive) Scale. Use 3 gallons per 100 gallons of water. Apply during dormant or delayed dormant period.

Pears: Clover (Brown Almond) Mite, European Red Mite. To control overwintering Mite eggs use 2 gallons per 100 gallons of water. Oystershell Scale, San Jose Scale, Pear Psylla. Use 2 gallons per 100 gallons of water. Apply during dormant or delayed dormant period.

Plums, Prunes: Clover (Brown Almond) Mite, European Red Mite, Leaf Curl Plum Aphid, Mealy Plum Aphid. To control overwintering Mite and Aphid eggs use 2 gallons per 100 gallons of water. Oystershell Scale, San Jose Scale. Use 2 gallons per 100 gallons of water. Paratonia (Olive) Scale. Use 3 gallons per 100 gallons of water. Apply during dormant or delayed dormant period.

Ornamentals: (Arborvitae, Ash, Birch, Boxwood, Catalpa, Dogwood, Elm, Euonymus, Juniper, Lilac, Magnolia, Maple, Myrica, Nandina, Oak, Pine, Poplar, Privet, Taxus, Tulip Tree, Willow, Yew, etc.) Use 2 gallons per 100 gallons of water. Cottony Maple, European Elm, European Spruce, Japanese Maple, and Tulip Tree Scale, Hard or Armored Scales, including Euonymus, Juniper, Oystershell, Pine Needle, Scaly, and Tulip Tree Scale. Deciduous ornamentals use 2 to 3 gallons per 100 gallons of water. Evergreen ornamentals use 2 gallons per 100 gallons of water. Apply as dormant spray after most danger of freezing nights has passed but before buds have begun to open. Spray on a clear, bright day and do not spray in late afternoon. Beware of frost damage. Do not use on easily injured plants such as Beech, Butternut, Hickory, Maple (Sugar & Japanese), or Walnut. Do not use on Douglas Fir flower buds and remove blue color from Colorado Blue spruce.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than that stated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and of handling and of storage of this material when such use and handling and of storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Time and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are left to the control of the seller and are assumed to be at the user's own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: FMC does not assume liability for damages for breach of warranty, or for consequential or special damages, or for expenses incurred, or for loss of profit, or for consequential damages.