

CODE
938

NET WEIGHT
50 LBS.

SEVIN* 10 SULFUR 40 DUST

FUNGICIDE - INSECTICIDE

DIRECTIONS FOR USE

COTTON: Boll weevil, bollworm, cotton fleahopper, lygus bugs, thrips, pink bollworm, cotton leaf worm, flea beetle, leaf perforator - Use 15 to 20 pounds of dust per acre. Apply when insects first threaten damage to crop and repeat at 5 to 7 day intervals as required to maintain control. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

RICE, FORAGE GRASS AND PASTURE: Thrips, armyworms, stink bug, grasshoppers - Use 10 to 15 lbs. of dust per acre. Apply when insects first threaten damage to crop. Repeat as necessary. Allow 14 days between last application and rice harvest. May be used within day of harvest or grazing of grass and pasture.

CORN: Corn earworm, European corn borer, fall armyworm, sap beetles, Japanese beetles - Use 10 to 20 lbs. of dust per acre. Apply when insects first threaten damage to crop. Repeat as necessary. For control of European corn borer, consult your state agricultural experiment station for proper timing of treatments. For corn earworm control, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Re-apply immediately after rain. May be used within day of harvest except when forage is fed to livestock then allow 7 days between last application and harvest. Application to the tassel region of corn during the pollen shed period will seriously reduce bee population.

SORGHUM (Grain and Forage): Armyworms, webworm, corn earworm, stink bug, grasshoppers - Use 10 to 20 lbs. of dust per acre. Apply when insects first threaten damage to crop. Repeat as necessary. Allow 20 days between last application and harvest of grain. No time limit on forage.

E. P. A. Reg. No. 279-1743

ACTIVE INGREDIENTS:

Sulfur ----- 40.0%

*Carbaryl 1(Naphthyl N-methyl-carbamate) ----- 10.0%

INERT INGREDIENTS: ----- 50.0%
100.0%

Sulfur fineness: 93% through 325 mesh.

*Sevin, U. S. Pat Nos. 2,903,478; 3,009,855

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

POTATOES: Lygus bugs, leafhoppers, fleabeetles - Use 20 lbs. per acre for lygus bugs, leafhoppers, and flea beetles. Apply when insects appear and repeat as necessary.

BEANS: Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetle, flea beetle, velvetbean caterpillar, corn earworm, army worm, stink bugs, bean rust - Use 10 to 15 lbs. of dust per acre. Apply when insects or diseases first appear. Repeat as required to maintain control. May be used up to day of harvest.

PEACHES: Peach twig borer, Oriental fruit moth, catfacing insects - Use 40 to 50 lbs. per acre. For control of Oriental fruit moth, apply as petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 1 day of harvest.

USAGE CAUTIONS

Applications of material containing sulfur should not be made just prior to or during periods of excessively high temperatures.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following dusting.

Do not mix with lime, Bordeaux, lime sulfur, casein-lime spreaders or other alkaline materials.

Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. ~~Keep out of smoke.~~

ACCEPTED

9-8-72

279-1743

Made in U.S.A.

