

DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary. May be applied with airplane or ground application equipment.

ON COTTON to control Aphids, Boll Weevil, Brown Cotton Leafworm, Cotton Fleahopper, Cotton Leafworm, Lygus Bugs, Mite, Thrips, Rapid Plant Bug, Tarnished Plant Bug and Stink Bug use 16-33 pounds per acre. Do not apply within 1 day of picking. If late season applications are made, do not graze livestock on treated areas or feed gin wastes.

ON APPLES, CRABAPPLES, PEARS and QUINCES to control Aphids, Apple Maggot, Codling Moth, European Apple Sawfly, Eye Spotted Bud Moth, Fruit Tree Leafroller, Leafhoppers, Mites and Red Banded Leafroller use 35-70 pounds per acre. Do not apply more than 8 times per season nor within 15 days of harvest.

ON CHERRIES to control Eye-spotted Bug Moth, Fruit Tree Leafroller, Mites and Plum Curculio use 35-80 pounds per acre. Do not apply more than 8 times per season nor apply within 15 days of harvest.

ON NECTARINES and PEACHES to control Aphids, Mites, Platynota Flavendana Leafroller and Red-banded Leafroller use 35-80 pounds per acre. Do not apply more than 8 times per season nor within 21 days of harvest.

ON PLUMS and PRUNES to control Aphids, Eye-Spotted Bud Moth, Fruit Tree Leafroller, Mites and Red-banded Leafroller use 35-80 pounds per acre. Do not apply more than 8 times per season nor apply within 15 days of harvest.

ON BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES and RASPBERRIES to control Aphids, Leafhoppers, Leafminers, Leafrollers, Obscure root weevil and Powdery Mildew apply 20-30 pounds per acre. Do not apply within 14 days of harvest.

ON GRAPES to control Grape Berry Moth, Leafhoppers, Mites, Thrips (California) and Powdery Mildew use 20-30 pounds per acre. To control Red banded Leafroller use 25-30 pounds per acre. Do not apply more than 3 applications per season nor within 28 days of harvest. Ground application preferred.

ON BROCCOLI, BRUSSELS SPROUTS, CABBAGE and CAULIFLOWER to control Aphids, Cabbage Looper, Diamondback Moth, and Imported Cabbageworm use 20-25 pounds per acre. Do not apply within 7 days of harvest for Brussels Sprouts, 15 days of harvest for broccoli and cauliflower, nor within 21 days of harvest for cabbage.

ON SPINACH to control Aphids, Leafminers and Mites use 15-25 pounds per acre. Do not apply within 14 days of harvest.

ON TOMATOES to control Colorado Potato Beetle, Green Stinkbug, Drosophila and Leafminers use 15-25 pounds per acre. To control Aphids, European Corn Borer, Flea Beetles, Hornworm, Leafhoppers and Thrips use 20-25 pounds per acre. To control Corn Earworm, Fruitworm, Pinworm, Tuberworm, and Yellow-striped Armyworm use 25 pounds per acre. No time limitations.

ON ORNAMENTALS and NURSERY PLANTS to control Aphids, Lacebugs, Leafhoppers, Mites and Thrips use 20 pounds per acre. Dust all foliage surfaces including the underside of leaves for complete coverage. Do not allow children or pets on treated areas until material has been washed into soil and treated area is dry.

CODE
31031

50 POUNDS
NET WEIGHT

SULPHUR 35 G-3 DUST

CONTAINS 3% GUTHION
INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-1827

ACTIVE INGREDIENTS:

BY WT.

0,0-Dimethyl S-(4-oxo-1,2,3-benzotriazin-3-ylmethyl)phosphorodithioate	3.00%
Sulphur	35.00%

INERT INGREDIENTS - - - - - 62.00%

U.S. Patent No. 2,758,115
Canadian Patent No. 552,660


100.00%

 **DANGER - POISON** 
KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.



Guthion is a registered trademark, Reg. U.S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G., Chemagro Corporation Licensee.

WARNING: Poisonous If Swallowed. Do not get in eyes or on skin. Wear rubber gloves, and goggles. Use respirator effective for protection to suitable types of masks or Department of Agriculture. Do not allow of reach of children. Keep all treated area or vicinity where treated who must enter treated fields wear protective clothing. Wash with soap and warm water before eating. Wash clothing with soap and hot water.
SYMPTOMS OF POISONING: A sense of constriction in the throat. Contracted pupils. Stomach pain.
TREATMENT: In case of poisoning, patient lie down and keep quiet. Give a tablespoonful of salt water or milk or water freely and again clear. If on skin, remove clothing immediately with soap and warm water with flowing water for at least 15 minutes.

USA

Do not graze livestock in treated areas. Do not treat food crops grown in treated areas. This product is toxic to fish and aquatic life in ponds, lakes, and streams. Apply this product carefully. **DISPOSE OF WASTES** by burying in soil or by burning. Keep Out of Smoke. Applications of materials containing Guthion should be made prior to or during periods of rain or in 10 days to two weeks of an application. Do not allow this material to be washed into water. This product is highly toxic to aquatic life. Residues on crops. Protective clothing. Cooperative Agricultural Extension Service.

assary. May be applied

own Cotton Leafworm,
Mite, Thrips. Rapid
use 16-33 pounds per
f late season appli-
reated areas or feed
ontrol Aphids, Apple
Eye Spotted Bud Moth,
Red Banded Leafroller
an 8 times per season

ruit Tree Leafroller,
acre. Do not apply
15 days of harvest.
s. Mites, Platynota
use 35-80 pounds per
on nor within 21 days

tted Bud Moth, Fruit
ler use 35-80 pounds
ason nor apply within

d RASPBERRIES to con-
trollers, Obscure root
er acre. Do not apply

ppers, Mites, Thrips
ds per acre. To con-
r acre. Do not apply
28 days of harvest.

ULIFLOWER to control
Imported Cabbageworm
7 days of harvest for
oli and cauliflower,

tes use 15-25 pounds
t.
le, Green Stinkbug,
er acre. To control
worm, Leafhoppers and
Corn Earworm, Fruit-
myworm use 25 pounds

hids, Lacebugs, Leaf-
re. Dust all foliage
complete coverage. Do
til material has been

CODE
31031

50 POUNDS
NET WEIGHT

SULPHUR 35 G-3 DUST

CONTAINS 3% GUTHION
INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-1827

ACTIVE INGREDIENTS:	BY WT.
* 0,0-Dimethyl S-(4-oxo-1,2,3-benzotriazin-3(4H)-ylmethyl)phosphorodithioate	3.00%
Sulphur	35.00%

INERT INGREDIENTS ----- 62.00%

† U.S. Patent No. 2,758,115
Canadian Patent No. 552,660

100.00%

 **DANGER - POISON** 
KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine. Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

ACCEPTED

Sep 15, 1969

UNITED STATES DEPARTMENT OF AGRICULTURE
WASHINGTON, D.C. 20250
FEDERAL AGRICULTURAL MECHANIZATION SERVICE
FARM LABOR DIVISION
WASHINGTON, D.C. 20250

Guthion is a registered trademark, Reg. U.S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G., Chemagro Corporation Licensee.

WARNING: Poisonous If Swallowed, Inhaled, or Absorbed Through Skin. Do not get in eyes or on skin. Wear protective clothing, natural rubber gloves, and goggles. Do not breathe dust. Wear a mask or respirator effective for protection against GUTHION. Information as to suitable types of masks or respirators is available from the U.S. Department of Agriculture. Do not contaminate feed or food. Keep out of reach of children. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Workers who must enter treated fields within 2 days of application should wear protective clothing. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

TREATMENT: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

USAGE CAUTION

Do not graze livestock in treated areas for 21 days after treatment. Do not treat food crops grown in the green house.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Apply this product only as specified on the label.


DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty containers should be disposed of by burying with wastes or by burning. Keep Out of Smoke.

Applications of materials containing Sulphur should not be made just prior to or during periods of excessively high temperature or within 10 days to two weeks of an oil spray.

Do not allow this material to drift onto sulfur sensitive crops

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE — On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY — Niagara warrants that this product, as it appears for sale by Niagara, conforms to the ingredient statement on the label, and it is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES — Buyer's sole and exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damage, not exceeding the purchase price paid, and shall not include incidental or consequential damages. Notice of any claims must be reported to Niagara within 90 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS — Because of certain uncontrollable conditions created or caused by force majeure or other which Niagara cannot control, buyer and user agree, on purchase or use, to assume responsibility for and release Niagara from, and to absolve Niagara of the warranty or use of the product attributable to such cause. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

