

SUMMER FLOWABLE EMULSION

LIGHT-MEDIUM

DIRECTIONS FOR MIXING

Before using this product, be sure the spray tank is clean and contains just enough water to cover the agitator shaft. With the agitator in motion (120 RPM or more), add the amount of spray oil required for a full tank. Pump this mixture through the overflow and back into the tank for a minute or two. The mixture should turn white, indicating good emulsification, then fill the tank with water. Keep agitator running at all times.

If an injector is used, first add 25 to 50 gallons of water to tank with engine running to provide good pressure, then put injector suction tube into the measured amount of spray oil, opening injector valve to allow the oil to be sucked into the tank. Fill the tank with water.

DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary. Apply the following recommended rates in sufficient water to thoroughly cover one acre.

ORANGES, LEMONS, GRAPEFRUIT: Yellow Scale, Citricola Scale, Red Scale, Brown Soft Scale — Use 1½ to 2 gallons of this material per 100 gallons of water.

APPLES, PEARS, PEACHES, PLUMS, PRUNES, CHERRIES, ALMONDS: Spider Mites — Use 1 to 1½ gallons per 100 gallons of water in delayed dormant or summer periods. Certain varieties of plums and prunes may be injured by oil sprays, check tolerance to oil prior to treatment.

OLIVES: Olive Scale, Black Scale, Oleander Scale — Use 1½ to 2 gallons of this material per 100 gallons of water. Apply in the post bloom through post harvest periods.

PEARS: After harvest for the control of Blister Mite, Bud Mite, Red Spider Mites — Use 1½ to 2 gallons in combination with 4 to 6 pounds of Kolofog per 100 gallons of dilute spray.

INSECTICIDE-MITICIDE

EPA REGISTRATION NO. 279-1861 AA

ACTIVE INGREDIENT:	BY WT.
Petroleum Oil	80.00%
INERT INGREDIENTS:	20.00%
	100.00%

Classification — Light Medium

Unsulphonated Residue — 92% Minimum

USAGE CAUTIONS

Use only in equipment with mechanical or paddle agitation. Use efficient equipment. Avoid application of oil sprays just prior to or during hot weather, rainy weather or drying winds. Apply when temperature is above 40° F. Do not apply just prior to freezing weather. Use only when trees are in healthy condition, and when moisture conditions are sufficient. Avoid over spraying.

Do not make an oil application within 2 weeks prior to or after an application of sulphur. On Citrus do not make an oil application within 60 days prior to or after an application of Sulphur.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Do Not Use if Material Does Not Emulsify. Do Not Use In Undiluted Form.

CAUTION

KEEP OUT OF REACH OF CHILDREN


Harmful if Swallowed. Avoid inhalation of spray mist.

For Emergency Assistance Call
716-735-3765

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Fish and other beneficial aquatic life may be harmed by this product. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not re-use empty 28 gallon drum. Return to drum reconditioner or destroy it by perforating or crushing and burying in a safe place away from water supplies.



054W

MANUFACTURED IN U.S.A. BY

FMC CORPORATION

AGRICULTURAL CHEMICAL DIVISION

MIDDLEPORT, N.Y.
FRESNO, CA.

YAKIMA, WA
RICHMOND, CA

FMC Chemicals

EPA EST., 279-CA-2 Printed in U.S.A.

279-1861
8-1-74
1081