


CODE
2070

NET WEIGHT
30 LBS.



THIRAM 65 WETTABLE POWDER

FUNGICIDE

DIRECTIONS FOR USE

Observe days interval between last application and harvest indicated in () following crop. Do not use in undiluted form.

APPLES (0): Scab — Use $1\frac{1}{2}$ to 2 pounds per 100 gallons of water. Apply in pre-bloom, calyx and early cover sprays. In later cover sprays, use 1 pound per 100 gallons of water as required to maintain control.

Cedar Apple Rust — Apply as directed for scab control. If recommended by your State Agricultural Extension Service or Experimental Station, apply a special bloom spray at $\frac{3}{4}$ pound per 100 gallons of water. Keep fruit and foliage covered during infection periods.

Apple Blotch, Bitter Rot, Black Pox, Botryosphaeria Spot, Brooks' Spot, Fly Speck, Sooty Blotch — Apply as directed for scab control. When weather conditions are conducive for disease development shorten interval between applications.

PEACHES (7): Brown Rot — Use $1\frac{1}{2}$ to 2 pounds per 100 gallons of water. For blossom blight, apply at pink stage and at 3 to 4 day intervals during bloom, timed to keep the newly opened blossoms covered. For fruit rot apply at petal-fall and first cover. Beginning one month before harvest apply at weekly intervals.

Rhizopus Rot — Apply as directed for fruit phase of brown rot control.

EPA REG. NO. 279-1872

ACTIVE INGREDIENT:

Thiram (tetramethylthiuram disulfide) 65.0%

INERT INGREDIENTS..... 35.0%
100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN.
For Emergency Assistance Call
716-735-3765

May cause skin irritation. Do not inhale dust or spray mist. Do not get in eyes or on skin. Wash thoroughly after handling. Consumption of alcoholic beverages increases the possibility of harm.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and **DISPOSE OF WASTES** by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

Scab — Apply at petal fall and repeat 3 to 6 times at 10 to 14 day intervals or until disease is under control. Use $1\frac{1}{2}$ to 2 pounds per 100 gallons.

STRAWBERRIES (0): Gray Mold (Botrytis) — Use 2 to $2\frac{1}{2}$ pounds per 100 gallons of water applying at rates up to 200 gallons per acre. Make 3 to 5 applications at 10 day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results. If applied within 3 days of picking, remove residues by washing.

Leaf Spot, Leaf Scorch, Leaf Blight — Use 2 to $2\frac{1}{2}$ pounds per 100 gallons of water applying at rates up to 200 gallons per acre. Begin applications at first signs of disease symptoms, and continue applications at monthly intervals during the spring and fall. Remove residues by washing if applications are made within 3 days of harvest.

CELERY (FOR USE IN PLANT BEDS ONLY): Early Blight (Cercospora), Rhizoctonia Damping-off, Root Rot — Use $1\frac{1}{2}$ pounds in 100 gallons of water. Apply approximately 15 gallons of spray per 1200 square feet of bed area. Increase dosage to 25 gallons as plants become larger. Begin spraying soon after plants emerge and repeat at 3 day intervals. Under severe disease conditions shorten intervals between treatments. Continue applications until plants are ready to set in field.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals