



Code 2070

Pounds Net Weight

Thiram 65 Wettable Powder

Fungicide

EPA Reg. No. 279-1872

EPA Est. 279-

Active Ingredient:

Thiram (tetramethylthiuram disulfide) 65.00%

Inert Ingredients: 35.00%
100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

For Emergency Assistance Call 716-735-3765.

See Other Panels for Additional Precautionary Information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution

May cause skin irritation. Do not inhale dust or spray mist. Do not get in eyes or on skin. Wash thoroughly after handling. Consumption of alcoholic beverages increases the possibility of harm.

Environmental Hazards

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State, or local disposal authorities for approved alternative procedures.

Observe days interval between last application and harvest indicated in () following crop. Do not use in undiluted form.

Apples (0): Scab—Use 1½ to 2 pounds per 100 gallons of water. Apply in prebloom, calyx and early cover sprays. In later cover sprays, use 1 pound per 100 gallons of water as required to maintain control. Cedar Apple Rust—Apply as directed for scab control. If recommended by your State Agricultural Extension Service or Experimental Station, apply a special bloom spray at ½ pound per 100 gallons of water. Keep fruit and foliage covered during infection periods. Apple Blotch, Bitter Rot, Black Pox, Botryosphaeria Spot, Brooks' Spot, Fly Speck, Sooty Blotch—Apply as directed for scab control. When weather conditions are conducive for disease development shorten interval between applications.

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103



Celery (for use in plant beds only): Early Blight (*Cercospora*) Rhizoctonia Damping-off, Root Rot—Use 1 1/2 pounds in 100 gallons of water. Apply approximately 15 gallons of spray per 1200 square feet of bed area. Increase dosage to 25 gallons as plants become larger. Begin spraying soon after plants emerge and repeat at 3 day intervals. Under severe disease conditions shorten intervals between treatments. Continue applications until plants are ready to set in field.

Peaches (7): Brown Rot—Use 1 1/2 to 2 pounds per 100 gallons of water. For blossom blight, apply at pink stage and at 3 to 4 day intervals during bloom, timed to keep the newly opened blossoms covered. For Fruit Rot apply at petal-fall and first cover. Beginning one month before harvest, apply at weekly intervals. Rhizopus Rot—Apply as directed for fruit phase of Brown Rot control. Scab—Use 1 1/2 to 2 pounds per 100 gallons. Apply at petal fall and repeat 3 to 6 times at 10 to 14 day intervals or until disease is under control.

Strawberries (8): Gray Mold (*Botrytis*)—Use 2 to 2 1/2 pounds per 100 gallons of water applying at rates up to 200 gallons per acre. Make 3 to 5 applications at 10 day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results. If applied within 3 days of picking, remove residues by washing. Leaf Spot, Leaf Scorch, Leaf Blight—Use 2 to 2 1/2 pounds per 100 gallons of water applying at rates up to 200 gallons per acre. Begin applications at first signs of disease symptoms, and continue applications at monthly intervals during the spring and fall. Remove residues by washing if applications are made within 3 days of harvest.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained. Use of this product in accordance with directions or established safe practices.

Damages: Buyer or user who is liable for damages or injury to property or persons shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental, consequential or other damages.

(279-10-60-PA)