

Code 2070

Pounds Net Weight

Thiram 65 Wettable Powder

Fungicide

EPA Reg. No. 279-1872

EPA Est., 279-

Active Ingredient:

Thiram (tetramethylthiuram disulfide) 65.00%

Inert Ingredients: 35.00%

100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution

May cause skin irritation. Do not inhale dust or spray mist. Do not get in eyes or on skin. Wash thoroughly after handling. Consumption of alcoholic beverages increases the possibility of harm.

Environmental Hazards

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not allow this material to drift onto neighboring crops, trees or other areas or use in a manner or at a time other than in accordance with label directions because of the potential for crop injury, illegal residues or other undesirable results may occur.

For Emergency Assistance Call 718-735-3765.

See Other Panels for Additional Precautionary Information.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION
Area treated with Thiram 65 Wettable Powder on (insert date). Do not enter without appropriate protective clothing until sprays have dried (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

Storage and Disposal

Pesticide Storage

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Contain spills. Call FMC Collect (718) 735-3765.


To contain spill, liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a leaking container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary and properly designated area as allowed by State or Federal authorities, by burning. If burned, stay out of smoke.



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

2381A
PA 8834

Observe days interval between last application and harvest indicated in () following crop. Do not use in undrained form.

Apples (0): Scab—Use 1 1/2 to 2 pounds per 100 gallons of water. Apply in prebloom, calyx and early cover sprays; in later cover sprays use 1 pound per 100 gallons of water as required to maintain control. Cedar Apple Rust—Apply as directed for scab control. If recommended by your State Agricultural Extension Service or Experimental Station apply a special bloom spray at 1/4 pound per 100 gallons of water. Keep fruit and foliage covered during infection periods. Apple Blotch, Bitter Rot, Black Pox, Botryosphaeria Spot, Brooks' Spot, Fly Speck, Sooty Blotch—Apply as directed for scab control. When weather conditions are conducive for disease development shorten interval between applications.

Celery (for use in plant beds only): Early Blight (Cercospora), Rhizoctonia Damping-off, Root Rot—Use 1 1/2 pounds in 100 gallons of water. Apply approximately 15 gallons of spray per 1200 square feet of bed area. Increase dosage to 25 gallons as plants become larger. Begin spraying soon after plants emerge and repeat at 3 day intervals. Under severe disease conditions shorten intervals between treatments. Continue applications until plants are ready to set in field.

Peaches (7): Brown Rot—Use 1 1/2 to 2 pounds per 100 gallons of water. For blossom blight, apply at pink stage and at 3 to 4 day intervals during bloom, timed to keep the newly opened blossoms covered. For Fruit Rot apply at petal fall and first cover. Beginning one month before harvest, apply at weekly intervals. Rhizopus Rot—Apply as directed for fruit phase of Brown Rot control. Scab—Use 1 1/2 to 2 pounds per 100 gallons. Apply at petal fall and repeat 3 to 6 times at 10 to 14 day intervals or until disease is under control.

Strawberries (0): Gray Mold (Botrytis)—Use 2 to 2 1/2 pounds per 100 gallons of water applying at rates up to 200 gallons per acre. Make 3 to 5 applications at 10 day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results. If applied within 3 days of picking, remove residues by washing. Leaf Spot, Leaf Scorch, Leaf Blight—Use 2 to 2 1/2 pounds per 100 gallons of water applying at rates up to 200 gallons per acre. Begin applications at first signs of disease symptoms and continue applications at monthly intervals during the spring and fall. Remove residues by washing if applications are made within 3 days of harvest.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use. The results to be obtained, if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damage, not exceeding the purchase price paid and shall not include incidental or consequential damages.