

Code 31030

Net Weight
Pounds

G-3 Dust Insecticide

For Agricultural or Commercial Use Only.


EPA Reg. No. 279-1945

EPA Est. 279-

Active Ingredients:	By Wt
*O, O'-Dimethyl S-[1[4-oxo-1,2,3-benzoxazin-3-(2H)-yl] methyl] phosphorodithioate	30%
Inert Ingredients:	97.0%
	100.0%

Contains 3% Guthion®
 *Guthion® is a registered trademark of the Parent Company of
 Farbenfabriken Bayer GmbH, Leverkusen
 Canadian Patent No. 552,660

KEEP OUT OF REACH OF CHILDREN



First Aid

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

If In Eyes: Flush eyes for 15 minutes with plenty of water. Call a physician immediately.

If On Skin Or Clothing: Remove contaminated clothing and shoes immediately. Bathe with soap and water. Call a physician immediately.

Antidote

Note To Physicians: Active ingredient is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidotal. Give 4 mg intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If cyanotic give artificial respiration. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

For Emergency Assistance Call 716-735-3765.
 See Other Panels for Additional Precautionary Information



FMC Corporation
 Agricultural Chemical Group
 2000 Market Street
 Philadelphia, Pennsylvania 19103

PRECAUTIONARY STATEMENTS Hazards To Humans (& Domestic Animals)

Danger

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash with soap and hot water before reuse. Wear natural rubber gloves, protective clothing and goggles. Wear a mask or pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II. Keep all unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered for at least two days.

Environmental Hazards

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture, or residue that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers or bury in a safe place. Do not reuse container. Consult Federal, State or local disposal authorities for approved alternative procedures.

Not for use or storage in or around the home
 Do not graze livestock in treated areas for 21 days after treatment
 Do not treat food crops grown in the greenhouse

Apply when pests first appear and repeat as necessary. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

Apples (7), Crabapples (7), Pears (7), Quinces (7): Aphids, Apple Maggot, Codling Moth, European Apple Sawfly, Eyespotted Bud Moth, Fruittree Leafroller, Leafhoppers, Mites, Redbanded Leafroller—Use 35-70 pounds per acre. Do not apply more than 8 times per season.

Apricots (21), Nectarines (21), Peaches (21): Aphids, Mites, Platynota Flavendana Leafroller, Redbanded Leafroller—Use 35-80 pounds per acre.

Blackberries (14), Boysenberries (14), Loganberries (14), Raspberries (14): Aphids, Leafhoppers, Leafminers, Leafrollers, Obscure Root Weevils—Use 20-30 pounds per acre.

Boysenberries—See Blackberries

Broccoli (15), Brussels Sprouts (7), Cabbage (21), Cauliflower (15): Aphids, Cabbage Looper, Diamondback Moth, Imported Cabbageworm—Use 20-25 pounds per acre.

Brussels Sprouts—See Broccoli

Cabbage—See Broccoli

ACCEPTED

OCT 16 1980

Federal Insecticide,
 Fungicide, and Rodenticide Act,
 24 CFR Part 151, for the pesticide
 registered under EPA Reg. No. 279-1945

BEST AVAILABLE COPY

19
 EPA Reg. No. 279-1945
 Page 1 of 2

16-33 pounds per acre. If late season applications are made, do not graze livestock on treated areas or feed gin wastes

Crabapples—See Apples

Grapes (10): Grape Berry Moth, Leafhoppers, Mites, Thrips (California) — Use 20-30 pounds per acre. **Redbanded Leafroller**—Use 25-30 pounds per acre. Do not apply more than 3 applications per season. Ground application equipment preferred.

Loganberries—See Blackberries

Nectarines—See Apricots

Ornamentals, Nursery Plants: Aphids, Lacebugs, Leafhoppers, Mites, Thrips — Use 20 pounds per acre. Dust all foliage surfaces including the underside of leaves for complete coverage. Do not allow children or pets on treated areas until material has been washed into soil and treated area is dry

Peaches—See Apricots

Pears—See Apples

Plums (15), Prunes (15): Aphids, Eyespotted Bud Moth, Fruittree Leafroller, Mites, Redbanded Leafroller—Use 35-80 pounds per acre. Do not apply more than 8 times per season

Prunes—See Plums

Quince—See Apples

Raspberries—See Blackberries

Spinach (14): Aphids, Leafminers, Mites — Use 15-25 pounds per season

Tomatoes (0): Colorado Potato Beetle, Green Stinkbug, Drosophila, Leafminers—Use 15-25 pounds per acre. Aphids, European Corn Borer, Flea Beetles, Hornworm, Leafhoppers, Thrips — Use 20-25 pounds per acre. Corn Earworm, Fruitworm, Pinworm, Tuberworm, Yellowstriped Armyworm—Use 25 pounds per acre.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use

On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

(31030—PA 6/80)

USA
[Stamp]

[Barcode]

1/2

EPA Reg. No. 279-1945

Code 31030

Net Weight
Pounds

G-3 Dust Insecticide

For Agricultural or Commercial Use Only.


EPA Reg. No. 279-1945

EPA Est. 279-

Active Ingredients:	By Wt
*O, O - Dimethyl S - [4-oxo-1, 2, 3 benzotriazin 3 (4H)-yl] methyl] phosphorodithioate	30%
Inert Ingredients:	97.0%
	100.0%

Contains 3% Guthion®
*Guthion® is a registered trademark of the Parent Company of
Farbenfabriken Bayer GmbH, Leverkusen
Canadian Patent No. 552,660

KEEP OUT OF REACH OF CHILDREN



First Aid

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

If in Eyes: Flush eyes for 15 minutes with plenty of water. Call a physician immediately.

If On Skin Or Clothing: Remove contaminated clothing and shoes immediately. Bathe with soap and water. Call a physician immediately.

Antidote

Note To Physicians: Active ingredient is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidotal. Give 4 mg intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If cyanotic give artificial respiration. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

For Emergency Assistance Call 716-735-3785.

See Other Panels for Additional Precautionary Information

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia, Pennsylvania 19103

PRECAUTIONARY STATEMENTS

Hazards To Humans (& Domestic Animals)

Danger

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorb skin. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap after handling and before eating or smoking. Remove contaminated clothing with soap and hot water before reuse. Wear natural rubber gloves, protective goggles. Wear a mask or pesticide respirator jointly approved by the Mining and Safety Administration (formerly the U.S. Bureau of Mines) and the National Occupational Safety and Health under the provisions of 30 CFR Part 11. Keep all persons and beneficial animals out of operating area or vicinity where there may be drift. Vacated areas should not be re-entered for at least two days.

Environmental Hazards

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residue. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open containers are prohibited. Pesticide, spray mixture, or rinseate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or in a safe place away from water supplies. Dispose of container in an incineration or landfill approved for pesticide containers or bury in a safe place. Do not reuse container. Consult Federal, State or local disposal authorities for appropriate procedures.

Not for use or storage in or around the home.
Do not graze livestock in treated areas for 21 days after treatment.
Do not treat food crops grown in the greenhouse.

Apply when pests first appear and repeat as necessary. May be applied with airplane application equipment. Observe days interval between last application and indicated by number in () following the crop.

Apples (7), Crabapples (7), Pears (7), Quinces (7): Aphids, Apple Maggot, Core European Apple Sawfly, Eyespotted Bud Moth, Fruittree Leafroller, Leafhopper, Redbanded Leafroller - Use 35-70 pounds per acre. Do not apply more than once per season.

Apricots (21), Nectarines (21), Peaches (21): Aphids, Mites, Platynota Flavendana, Redbanded Leafroller - Use 35-80 pounds per acre.

Blackberries (14), Boysenberries (14), Loganberries (14), Raspberries (14): Aphids, Hoppers, Leafminers, Leafrollers, Obscure Root Weevils - Use 20-30 pounds per acre.

Boysenberries - See Blackberries.

Broccoli (15), Brussels Sprouts (7), Cabbage (21), Cauliflower (15): Aphid, Looper, Diamondback Moth, Imported Cabbageworm - Use 20-25 pounds per acre.

Brussels Sprouts - See Broccoli.

Cabbage - See Broccoli.

Page 2 of 2 779. 1965

Cauliflower—See Broccoli

Cherries (15): Eyespotted Bud Moth, Fruitree Leafroller, Mites, Plum Curculio—Use 35-80 pounds per acre. Do not apply more than 8 times per season

Cotton (1): Aphids, Boll Weevil, Brown Cotton Leafworm, Cotton Fleahopper, Cotton Leafworm, Lygus Bugs, Mites, Thrips, Rapid Plant Bug, Tarnished Plant Bug, Stink Bug—Use 16-33 pounds per acre. If late season applications are made, do not graze livestock on treated areas or feed gin wastes

Crabapples—See Apples

Grapes (10): Grape Berry Moth, Leafhoppers, Mites, Thrips (California)—Use 20-30 pounds per acre. Redbanded Leafroller—Use 25-30 pounds per acre. Do not apply more than 3 applications per season. Ground application equipment preferred

Loganberries—See Blackberries

Nectarines—See Apricots

Ornamentals, Nursery Plants: Aphids, Lacebugs, Leafhoppers, Mites, Thrips—Use 20 pounds per acre. Dust all foliage surfaces including the underside of leaves for complete coverage. Do not allow children or pets on treated areas until material has been washed into soil and treated area is dry

Peaches—See Apricots

Pears—See Apples

Plums (15), Prunes (15): Aphids, Eyespotted Bud Moth, Fruitree Leafroller, Mites, Redbanded Leafroller—Use 35-80 pounds per acre. Do not apply more than 8 times per season.

Prunes—See Plums

Quince—See Apples

Raspberries—See Blackberries

Spinach (14): Aphids, Leafminers, Mites—Use 15-25 pounds per season

Tomatoes (8): Colorado Potato Beetle, Green Stinkbug, Drosophila, Leafminers—Use 15-25 pounds per acre. Aphids, European Corn Borer, Flea Beetles, Hornworm, Leafhoppers, Thrips—Use 20-25 pounds per acre. Corn Earworm, Fruitworm, Pinworm, Tuberworm, Yellowstriped Armyworm—Use 25 pounds per acre

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use

On purchase of this product buyer and user agree to the following conditions

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages