CODE 2258





279-158.

SPREADERSUTTINGScott#4Lawn Beauty#5

#4

Approximate fertilizer spreader settings to distribute 4 lbs. of Niagara Section 59

USAGE CAUTIONS

Granular per 1000 sq. ft.

NULAR

Do not apply on Boston Ivy, as injury may occur.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGH-BORING CROPS OR NON-CROP AREAS OR USE IN A MAN-NER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UN-DESIRABLE RESULTS MAY OCCUR.

Dealers Should Seil in Original Packages Only.

TERMS OF SALE OR USE On purchase or use of this product buyer and user agree to the following conditions

WARRANTY Niagara warrants that this product, as of the birsh of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set both it, the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED

DAMAGES Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited that drives damages not exceeding the purchase price paid and shall be include incidental or consequential damages. Notice of any claim must be reported to Niggara within 10 days of discourt, by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS Because of certain critical conditions created or incurrent by house er or user or over which Niagara has no contrôl, huver and user, as a condition of purchase or use, assume responsibility for and release Niagara from, Itability arising out of the handling or use of the product attributable to such causes. No person us authorized to vary or waive any statement contained herein

DIRECTIONS FOR USE

SEVIN*


<u>ON CORN</u> to control European Corn Borer use 30-35 pounds per acre, for first generation European Corn Borer larvae in whorls. Use suitable ground or air equipment to insure good coverage. Apply when pests appear in damaging numbers and repeat 7 to 14 days later if necessary.

ON LAWN to control Chinch Bugs, Sod Webworms or Lawn Moths, Millipeds, Earwigs, Fleas, Ants, Leafhopper and Fall Armyworms, use 4 pounds per 1000 sq. ft. treating the lawn evenly. For best results cut the lawn, remove clippings and water before making a treatment. Work material in with a coarse water spray after treatment.

Application of this material should be made at the first signs of insect activity, and repeat applications should be made as necessary to maintain effective control.

25 POUNDS COVERS 6000 SQUARE FEET.

For use by Pest Control Operators only to control Darkling Ground Baetle, Earwigs and Brown Dog ticks. Use this material at the rate of 1 pound per 250 square feet. Spread this material in the infested areas at the first sign of the pests.



CONTAINS 5% SEVIN INSECTICIDE

INSECTICIDE

USDA REG. NO. 279-1991

ACTIVE INGREDIENT:

*Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS:	95.0%
	00.0%

*Sevin, U. S. Pat. No. 2,903,478; 3,009,855

CAUTION KEEP OUT OF REACH OF CHILDREN


Harmful if inhaled or swallowed. Avoid prolonged or repeated breathing of dust. Do not take internally. Skin contact may be harmful. Avoid prolonged or repeated contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOT E FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following dusting.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

MANUFACTURED IN U.S.A. BY



*Trademark of Union Carbide Corporation for the active ingredient Carbaryl,

017-R\$ 8. 150