

Code: 30899

50 Pounds

# KELTHANE\* SULPHUR 4-25 DUST

## DIRECTIONS FOR USE

Apply before mites build up to the point where they are damaging the crop or when disease first appears and repeat as necessary. For disease control this material must be applied prior to disease development. For best results, all portions of the plant must be covered.

For disease control after the time limitations of this material use a suitable material.

May be applied with ground or airplane application equipment.

**ON GRAPES, RASPBERRIES, BLACKBERRIES, DEWBERRIES, LOGANBERRIES and ROYSENBERRIES** for the control of Two-Spotted, Pacific, and Willamette Mites and Powdery Mildew, use 30 pounds per acre. On Grapes apply 30-35 pounds per acre. Do not apply within two days of harvest on Raspberries and other cane berries or within 7 days of harvest on grapes. Use ground application equipment.

**ON STRAWBERRIES** for the control of Two-Spotted and Strawberry Mites, and Powdery Mildew use 30-60 pounds per acre. Do not apply within 2 days of harvest.

**ON BEANS** for the control of Powdery Mildew, Rust, and Spotted and Pacific Mites use 30-35 pounds per acre. Do not apply within 7 days of harvest. Do not feed treated vines to meat or dairy animals. Sulphur may injure some varieties, check tolerance before use.

**ON COTTON** for the control of Two-Spotted, Pacific, Atlantic and Desert Mites use 30-40 pounds per acre. Do not apply within 14 days of harvest. Do not feed treated forage to meat or dairy animals. Do not graze meat or dairy animals on treated areas.

**ON PEPPERS and TOMATOES** for the control of Two-Spotted, Pacific, Tropical and Tomato Russet Mites use 30-35 pounds per acre. Do not apply within 2 days of harvest.

**ON CHERRIES, NECTARINES and PEACHES** for the control of Mites, use 40-80 pounds per acre. Do not apply within 7 days of harvest on Cherries, or within 14 days of harvest on Nectarines or Peaches. Do not repeat applications on Cherries within 30 days of the preceding application.

**ON APPLES, CRABAPPLES and QUINCE** for the control of European Red, Clover, Two-Spotted, Schoene, McDaniel, Willamette, Pacific and Apple Rust Mites use 40-100 pounds per acre. Sulphur may injure some varieties, check sulphur tolerance prior to use. Do not apply within 7 days of harvest.

**ON PLUMS, PRUNES and FIGS** to control Mites, use 40-50 pounds per acre. Do not apply within 7 days of harvest. Do not reapply within 30 days of preceding application.

FUNGICIDE - MITICIDE

U.S.D.A. REGISTRATION NO. 279-1996

| ACTIVE INGREDIENTS:                               | BY WT.  |
|---|---------|
| 1,1-Bis (p-chlorophenyl) - 2,2,2-trichloroethanol | 4.00%   |
| Sulphur   | 25.00%  |
| INERT INGREDIENTS                                 | 71.00%  |
|   | 100.00% |

\*Trademark Rohm & Haas Co.

**ON ORNAMENTALS, NURSERY STOCK,** and Powdery Mildew and Mite. Dust for tolerances prior to treating any **ON CITRUS** (Oranges, Grapefruit, Lo Kumquats) to control Citrus Red-spotted and Yuma Mites apply 50-100 lbs within 7 days of harvest.

## USAGE CAUTION

Application of material's contained prior to or during periods of excessive rain. Do not allow this material to drift. This product is toxic to fish. Keep away from water. Apply only as specified on the label. **DISPOSE OF WASTES** by pouring in water. Empty bags should be burned. Out of Smoke.

## CAUTION


### KEEP OUT OF REACH OF CHILDREN

Avoid breathing dust

Avoid storage near feed and food products

Wash thoroughly after handling

Keep away from domestic animals, feed and foodstuffs



# 50 Pounds Net Weight

# KELTHANE\* SULPHUR 4-25 DUST

**FOR USE**

point where they are damaging the  
rs and repeat as necessary. For  
ust be applied prior to disease  
ll portions of the plant must be

limitations of this material use

airplane application equipment.

**LES, DEWBERRIES, LOGANBERRIES** and  
Two-Spotted, Pacific, and Willa-  
se 30 pounds per acre. On Grapes  
t apply within two days of harvest  
ies or within 7 days of harvest on  
ipment only.

Two-Spotted and Strawberry Mites,  
s per acre.  
est.

ry Mildew, Rust, Two-Spotted and  
acre. Do not apply within 7 days  
vines to meat or dairy animals.  
s, check tolerance before using.

o-Spotted, Pacific, Atlantic and  
acre. Do not apply within 14 days  
orage to meat or dairy animals. Do  
n treated areas.

control of Two-Spotted, Pacific,  
use 30-35 pounds per acre. Do not

HES for the control of Mites, use  
apply within 7 days of harvest on  
rvest on Nectarines or Peaches. Do  
s within 30 days of the preceding

for the control of European Red,  
Daniel, Willamette, Pacific and  
per acre. Sulphur may injure some  
prior to use. Do not apply within

ontrol Mites, use 40-50 pounds per  
of harvest. Do not reapply within

**FUNGICIDE - MITICIDE**

U.S.D.A. REGISTRATION NO. 279-1996

| ACTIVE INGREDIENTS:                | BY WT.         |
|------------------------------------|----------------|
| 1,1-Bis (p-chlorophenyl) - 2,2,2-  |                |
| trichloroethanol - - - - -         | 4.00%          |
| Sulphur - - - - -                  | 25.00%         |
| <b>INERT INGREDIENTS - - - - -</b> | <b>71.00%</b>  |
|                                    | <b>100.00%</b> |

\*Trademark Rohm & Haas Co.

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

- Avoid breathing dust
- Avoid storage near feed and food products
- Wash thoroughly after handling
- Keep away from domestic animals, feed and foodstuffs.

**ON ORNAMENTALS, NURSERY STOCK, and SHADE TREES** to control Rust, Powdery Mildew and Mite. Dust for complete coverage. Check sulphur tolerances prior to treating any plant.


**ON CITRUS** (Oranges, Grapefruit, Lemons, Limes, Tangeloes, Tangerines, Kumquats) to control Citrus Red, Citrus Flat, Citrus Rust, Six-spotted and Yuma Mites apply 50-100 pounds per acre. Do not apply within 7 days of harvest.

PAGE CAUTION


Application of material containing sulphur should not be made prior to or during periods of excessively high temperature.

Do not allow this material to drift to sulfur sensitive plants. This product is toxic to fish. Keep out of use, food, and streams. Apply only as specified on the label.

**DISPOSE OF WASTES** by burying in approved landfills away from water supplies. Empty bags should be covered with waste, or incinerated. Keep Out of Smoke.



**ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.**



**Dealers Should Sell in Original Packages Only.**

**DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.**

**TERMS OF SALE OR USE** On purchase or use of this product buyer and user agree to the following conditions

**WARRANTY** Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

**DAMAGES** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

**EXCUSE FROM NON-CONTROLLABLE CONDITIONS** Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



**NIAGARA CHEMICAL DIVISION**  
**FMC CORPORATION**  
**MIDDLEPORT, NEW YORK**

