

Code 2347

as amended, for the purpose of  
registered under  
EPA Reg. No.

Pounds Net Weight

# G-3 Dust

## Insecticide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-2022 EPA Est.

Active Ingredient:	By Wt
0,0-Dimethyl S-[4-oxo-1,2,3-benzotriazin-5 (4H)-ylmethyl] phosphorodithioate .....	3.0%
Inert Ingredients: .....	97.0%
	100.0%

Contains 3% Guthion\*

\*Guthion is Reg. TM in U.S. by Farbenfabriken Bayer, A.G. Chemagro Corp. Licensee

**KEEP OUT OF REACH OF CHILDREN**

**☠ DANGER—POISON ☠**

**First Aid:**

**If Swallowed:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

**If in Eye:** Flush eye for 15 minutes with plenty of water. Call a physician immediately.

**If on Skin or Clothing:** Remove contaminated clothing and shoes immediately. Bathe with soap and water. Call a physician immediately.

For Emergency Assistance Call 716-735-3765.

See Other Panels for Additional Precautionary Information.

# FMC

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

PA 6.80

**Antidote**

**Note to Physician:** Guthion is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidotal. Give 4 mg intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If cyanotic give artificial respiration. 2-PAM is also antidotal and may be administered in conjunction with atropine.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans (& Domestic Animals)

**Danger**

Poisonous if swallowed, inhaled or absorbed through skin. **Rapidly absorbed through skin.** Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11. Keep all unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered for at least two days. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not use on other crops used for feed or forage. Treated orchards should not be pastured.

#### Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or material that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not reuse this container. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures.

Not for use or storage in or around the home

Apply when insects first appear and repeat as required to maintain control. Observe days interval between last application and harvest indicated by the number in ( ) following the crop. Observe use limitations.

**Alfalfa (21), Clover (21):** Alfalfa Plant Bug, Aphids, Fleahoppers, Leafhoppers, Lygus Bugs, Mites, Spittlebug—Use 20 to 25 pounds per acre. Do not apply more than once between cuttings.

**Apples (7), Crabapples (7), Pears (7), Quinces (7):** Aphids, Apple Maggot, Codling Moth, European Apple Sawfly, Eyespotted Bud Moth, Fruitree Leafroller, Leafhoppers, Mites, Plum Curculio, Red-banded Leafroller—Use 35 to 40 lbs per acre. Do not apply more than 8 times per season.

**Apricots (21), Nectarines (21), Peaches (21):** Aphids, Mites, Plum Curculio, Redbanded Leafroller—Use 35 to 40 lbs per acre.

**Blueberries:** Blueberry Maggot, Lecanium Scale, Plum Curculio, Fruitworms, Leafminers—Use 20 lbs per acre. Do not apply more than 5 times per season. Do not apply within 9 hours of harvest.

**Broccoli (15), Brussels Sprouts (7), Cabbage (21), Cauliflower (15):** Aphids, Cabbage Looper, Imported Cabbageworm, Diamondback Moth Larvae—Use 20 to 25 lbs per acre.

**Cherries (15):** Eyespotted Bud Moth, Fruitree Leafroller, Mites, Plum Curculio—Use 35 to 40 lbs per acre. Do not apply more than 8 times per season.

**Grapes (10):** Grape Berry Moth, Leafhoppers, Mites, Redbanded Leafroller—Use 25 to 30 lbs per acre. Do not apply more than 3 times per season.

**Pine Trees:** European Pine Shoot Moth, Nantucket Pine Tip Moth—Use 15 to 30 lbs per acre. Use maximum rates for tip moth control in shade.

**Plums (15), Prunes (15):** Aphids, Eyespotted Bud Moth, Fruitree Leafroller, Mites, Plum Curculio, Redbanded Leafroller—Use 35 to 40 lbs per acre. Do not apply more than 8 times per season.

**Potatoes (7):** Aphids, Colorado Potato Beetle, Flea Beetle, Leafhoppers, Leafminers, Spittlebug, Tarnished Plant Bug—Use 20 to 25 pounds per acre.

**Tobacco (6):** Aphids, Tobacco Hornworm, Tobacco Flea Beetle—Use 20 to 25 lbs per acre. Treat for Hornworm as soon as egg masses or worms are first found. Second brood Hornworms may require two applications, one in late July and another in August. Flea beetles may require two applications, one about August first and another about two weeks later. Tobacco Budworm—Use 25 lbs per acre. Make application directly into bud. Repeat as necessary. Prime before treating.

### Dealers Should Sell in Original Packages Only.

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions.

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty, or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

(2347-6/80-PA)