

Code 141


Gals. Net Contents

Aqua Ethion 8


Insecticide
For Agricultural or Commercial Use Only.

EPA Reg. No. 279-2102 EPA Est.

Active Ingredients:	B. Wt
Ethion (0, 0, 0', 0'-Tetraethyl S, S -methylene bisphosphorodithioate)	81.0%
Inert Ingredients:	19.0%
Contains 8 lbs. Ethion per gallon	100.00%



KEEP OUT OF REACH OF CHILDREN



PRECAUTIONARY STATEMENTS
Hazards To Humans (& Domestic Animals)

First Aid Treatment

If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If on skin or clothing, immediately flush skin or eyes with water for at least 15 minutes while removing contaminated clothing and shoes.

If in eye, flush eye for 15 minutes with plenty of water. Call a physician immediately.

If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Note to Physician: Active ingredient is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidotal. Give 4 mg atropine sulfate intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. Do not give morphine.

See Other Panels for Additional Precautionary Information.
For Emergency Assistance Call 716-735-3765.



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PAGE

Danger

Extremely poisonous if swallowed, inhaled, or absorbed through skin. Do not breathe vapor or spray mist. Do not get in eyes, on skin or on clothing. In case of contact, wash immediately with water. For eyes flush with plenty of water and get medical attention. Wear a mask or respirator of a type approved by the Mining Enforcement and Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11 for Ethion protection. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

Chestnut, Filbert, Pecan (0): Aphids, Mites—2 1/2 pints per acre. Do not apply more than twice during the fruiting period. On chestnuts and pecans, do not apply after husk split. On filberts, do not apply when nut is visible. Do not graze livestock in treated groves.

Corn, Field (50): Pacific Mite, Twospotted Spider Mite—1 pint per acre. Do not make more than 1 application after ear formation. Do not feed treated fodder to livestock.

Cotton: Aphids—1 to 1 1/2 pt per acre. Spider Mites—1 to 1 1/2 pt per acre. Cotton Leafworm—1 pint per acre. Lygus Bugs, Spider Mites (W. E. COAST AREA ONLY)—1 to 1 1/2 pints per acre. Treat when an infestation reaches the point where control is required and repeat as necessary. Do not apply after the bolls open. Do not graze dairy or meat animals in treated fields.

Cucumber (0), Summer Squash (0): European Leafminer, Twospotted Spider Mite—1 pt per acre.

Eggplant (0): Twospotted Spider Mite—1 to 1 1/2 pt per acre.

Melon (0): Melon Leafminer, Serpentine Leafminer, Twospotted Spider Mite—1 pt per acre. Melon Leafhopper—1 to 1 1/2 pt per acre. Petrosia Mite (California only)—1 to 1 1/2 pints per acre.

Nectarine (30): Clover Mite, European Red Mite, Twospotted Spider Mite—1 pint per 100 gals. or 1 to 1 1/2 pints per acre. Do not exceed 2 applications during fruiting period.

Ornamentals, Greenhouse: Commercial—Mites, Aphids. TREATMENT: Use 5 fl. ozs. Add-Ethion in 15 gals. of water and apply to 2,500 sq. ft. of heated space. Wear mask and goggles. Ethion is protected by the Mine, Health, and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health. See the product label for CFR Part 16 when applying in greenhouse.

Ornamentals, Out-of-Doors: Commercial—Mites, Aphids, Spider Mite—1 pt per 100 gals. or 1 to 1 1/2 pints per acre. Do not exceed 2 applications per year following application of this material.

Pear (30): Clover Mite, European Red Mite, Twospotted Spider Mite—1 pint to 1 1/2 pints per 100 gals. Pear Psylla—1 to 1 1/2 pints per 100 gals. Do not exceed 2 applications after first cover spray.

Pepper (0), Pimento (0): European Leafminer, Twospotted Spider Mite—1 to 1 1/2 pt per acre.

Strawberry (2): Strawberry Leafminer, Twospotted Spider Mite—1 pt per 100 gals. or 1 to 1 1/2 pints per acre. Do not exceed 2 applications after harvest is required.

Sorghum (Grain & Forage): European Leafminer, Mites, Aphids, 1 to 1 1/2 pt per acre. Do not exceed 2 applications per year. Do not feed treated fodder to livestock. Do not graze dairy or meat animals in treated fields. Do not apply after the grain is ready for harvest.

Tomato (2): European Leafminer, Twospotted Spider Mite—1 to 1 1/2 pt per acre.

Turf: Commercial Use: Control Fleas, Sod Webworm—1 to 1 1/2 pt per 100 sq. ft. of lawn applied in 10 gals. of water or 1 to 1 1/2 pints in 100 gals. of water to treat 500 sq. ft. To treat one acre, use 3 to 5 qts. applied in 400 gals. of water. It is important that the spray penetrates through the mat of turf to reach the insects. It helps to have the lawn well irrigated prior to treatment.

CAUTION: Keep children and animals off treated areas until this material has been thoroughly watered into soil and the grass has completely dried.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product, buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and of handling and of storage of this material when such use and of handling and of storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Time and method of application, weather and crop conditions, mixture with other chemicals, not specifically recommended, and other influencing factors, the use of this product are beyond the control of the seller and a responsibility of the user.

Use of Product: FMC's recommendations for the use of this product are based on tests believed to be reliable. The use of this product is not a guarantee of the control of the manufacturer, no guarantee is expressed or implied, is made as to the effects of such use, or the results obtained, if not used in accordance with directions or established practices.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

151 PA 6 60



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not store at temperatures below 20° F. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. For container disposal: triple rinse (or equivalent) or reuse and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place. Consult Federal, State, or local disposal authorities for approved alternative procedures. Do not reuse container. Do not cut or weld this container.

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Where amount of water is not specified, use with sufficient water for thorough coverage. For aircraft application to field crops and vegetables, mix the recommended amounts with sufficient water to provide a MINIMUM of 1 gallon of finished spray per acre. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in parentheses: (1) following crop. Do not use in undiluted form.

Apple (60): Codling Moth: 1/2 pint per 100 gals. Do not exceed 4 applications during fruiting period.

Apple (40): Codling Moth: 1/2 pint per 100 gals. Do not exceed 2 applications during fruiting period. Clover Mite, European Red Mite, Twospotted Spider Mite: 1/2 pint per 100 gals. Do not exceed 4 applications during fruiting period.

Apple (20): Clover Mite, European Red Mite, Twospotted Spider Mite: 1/2 pint per 100 gals. Do not exceed 2 applications during fruiting period.

Do not use on Astrachen, Davey, Duchess, Margaret Pratt, Melba, Wealthy, Williams Red (Red Williams), Yellow Transparent or on Crimson and N.J. #3 (Brite Mac), or any varieties maturing with or before Early McIntosh, as injury is likely to occur.

Beans (Succulent and Dry) (2): Bean Leaf Beetle, Mexican Bean Beetle, Twospotted Spider Mite: 1/2 to 1 pt. per acre. Flat Leafed Beet: 1/2 pt. per acre. Do not feed bean foliage to livestock.

Beans (Succulent and Dry) (4): Lygus Bug: 1/2 pt. per acre. Do not feed treated bean foliage to livestock.

Bermuda Grass (Lawn and Field Crop): Eriophyid Mites: 1/2 tablespoon per gallon of water to cover 200 sq. ft. of lawn. Use 1/2 ozs. per 100 gals. of water per 200 sq. ft. Applications needed to maintain effective control. Keep children and animals off treated areas until the spray has completely dried. Do not apply to Bermuda grass to be used for grazing or forage to be fed to livestock.