

CODE
2424

NET WEIGHT
50 LBS.

G-2 DUST

INSECTICIDE

DIRECTIONS FOR USE

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

Begin applications when pests first appear and repeat applications as needed to maintain control. Thorough coverage is essential for good control. Observe days interval between last application and harvest as indicated by number in parenthesis following crop.

BEANS, SNAP (7): Aphids, Leafhoppers, Mexican Bean Beetles, Mites. Use 30 lbs. dust per acre. Do not feed or ensile treated forage.

BLACKBERRIES (14): Aphids, Leafhoppers, Leaf Miners, Leaf Rollers. Use 25 lbs. dust per acre.

BLUEBERRIES: Blueberry Maggot, Fruitworm, European Scale, Plant Curculio. Use 30 lbs. dust per acre. Do not apply more than 8 times per season. Do not apply within 90 hours of harvest.


BOYSENBERRIES (14): Aphids, Leafhoppers, Leaf Miners, Leaf Rollers. Use 25 lbs. dust per acre.

BROCCOLI (15): Aphids, Cabbage Looper, Diamond Back Moth, Imported Cabbageworm. Use 25 lbs. dust per acre.

BRUSSELS SPROUTS: Aphids, Cabbage Looper, Diamond Back Moth, Imported Cabbageworm. Use 25 lbs. dust per acre.

CABBAGE (15): Aphids, Cabbage Looper, Diamond Back Moth, Imported Cabbageworm. Use 25 lbs. dust per acre.

CAULIFLOWER (15): Aphids, Cabbage Looper, Diamond Back Moth, Imported Cabbageworm. Use 25 lbs. dust per acre.



CONTAINS 2% GUTHION

EPA Reg. No. 279-2111

ACTIVE INGREDIENT:

10,0-Dimethyl S - 1-Ethoxy-1,2,3,4-tetrahydro-2,4-diazotriazin-3-one (4H)-ylmethyl phosphorothioate 98.0

INERT INGREDIENTS: 2.0

100.0

Guthion is Reg. TM of Farbwerke Bayer
A. G. Chem. Corp., Leverkusen,
U. S. Patent No. 2,758,115
Canadian Patent No. 855,700

WARNING

KEEP OUT OF REACH OF CHILDREN.

For Emergency Assistance Call

710-725-3765

May be fatal if swallowed, inhaled, or absorbed through skin. Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing, and cap. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Guthion protection. Keep all unprotected persons out of operating area or vicinity where there may be danger of drift. Vacated areas should not be reentered for at least two days. Do not store near food and food products. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

NOTE TO PHYSICIANS: Administer antidote 1-PAM (2-PAM) immediately and may require intubation and respiration.

DIRECTIONS FOR USE (CONT.)

CRANBERRIES (21): Tipworm. Use 50 lbs. dust per acre.

GRAPES (0): Flea Beetle, Grape Berry Moth, Leafhoppers, Leaf Rollers, Mites, Red-banded Leaf Roller. Use 35 lbs. dust per acre. Do not apply more than 3 times per season.

LOGANBERRIES (14): Aphids, Leafhoppers, Leaf Miners, Leaf Rollers. Use 25 lbs. dust per acre.

POTATOES (7): Aphids, Colorado Potato Beetle, Flea Beetle, Leafhoppers, Spittlebug, Tarnished Plant Bug, Tuber Moth. Use 35 lbs. dust per acre.

RASPBERRIES (14): Aphids, Leafhoppers, Leaf Miners, Leaf Rollers. Use 25 lbs. dust per acre.

PINACH (14): Aphids, Leafhoppers, Mites. Use 25 lbs. dust per acre.

TOBACCO (0): Aphids, Tobacco Budworm, Tobacco Hornworm, Tobacco Flea Beetle. Use 35 lbs. dust per acre. To achieve good Tobacco Budworm control, dust must be applied directly into bud. To achieve good Tobacco Hornworm control, apply dust as soon as egg masses or worms are first found. Do not dust after treating.

TOMATOES (0): Fleaworm, Hornworm, Leafminer. Use 35 lbs. dust per acre.

USAGE CAUTIONS

Do not use this product on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use.

Do not use this product on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use.

Do not use this product on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use. Do not use on or near food crops unless specifically labeled for such use.

WILSON JOURNAL

LABEL

Camerman's Note
Poor Copy

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT TO THE RIGHT. IT SHOULD BE KEPT IN THE ORIGINAL PACKAGE OR IN A CONTAINER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS. SEE ABOVE. AVOID ALL OTHER DIRECTIONS FOR USE OF THIS PRODUCT.

TERMS OF SALE: (1) On purchase or use of this product, the user must follow the following conditions:

(1) READ THE LABEL DIRECTIONS CAREFULLY. DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED. DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED. DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED.

(2) DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED. DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED. DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED.

(3) DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED. DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED. DO NOT USE THE PRODUCT IN ANY MANNER OTHER THAN AS DIRECTED.

MANUFACTURED IN U.S.A. BY

WILSON JOURNAL
1000 1000 1000
1000 1000 1000

1000 1000 1000