

[279-24]

FIRST AID

Call A Physician Immediately. **If Swallowed** Give 1 teaspoon of table salt in a glass of water. Repeat until vomiting occurs and vomitus is clear. Never induce vomiting in an unconscious person. **If In Eye** Flush eye for 15 minutes with plenty of water. **If On Skin or Clothing** Remove contaminated clothing and shoes immediately. Bathe with soap and water.

USAGE CAUTIONS

This product is toxic to fish and wildlife. Birds feeding in treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Apply this product only as specified on the label.

Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues, or other undesirable results may occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Store in Cool, Dry Place.

Not for use or storage in or around the home.

DISPOSAL WARNING: Do not re-use empty 2 and 5 gallon containers. Destroy them by perforating or crushing. Return drum to reconditioner or destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

**For Emergency Assistance Call
716-735-3765**


093W

Manufactured in U.S.A. By

MC Corporation

Agricultural Chemical Division

MIDDLEPORT, N.Y. YAKIMA, WA
FRESNO, CA RICHMOND, CA



**CODE
2486**

**NET CONTENTS
GALLONS**

PHOSPHAMIDON 8 SPRAY

INSECTICIDE

E.P.A. REGISTRATION NO. 279-2114 ZA

ACTIVE INGREDIENTS:	BY WT.
Phosphamidon (2-Chloro-2-diethylcarbamoyl-1-methylvinyl dimethyl phosphate)	75.50%
Related Compounds	2.50%
INERT INGREDIENTS:	22.00%
Contains 8 lbs. Phosphamidon Insecticide per Gallon	100.00%
U.S. Patent No. 2,908,601	

DANGER - POISON
KEEP OUT OF REACH OF CHILDREN

DANGER: Poisonous If Swallowed, Inhaled or Absorbed Through Skin! Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for phosphamidon protection. Rapidly Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs.

ANTIDOTE: NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidotal. Give 4 mg. intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If cyanotic give artificial respiration. Pralidoxime chloride (2-PAM) should be used in treatment.

READ ENTIRE LABEL AND DIRECTIONS, AND DOSAGE MAY WE RECOMMEND RE USAGE.

WARNING: Do not occur. Avoid drift to Phosphamidon 8 Sp and chewing insects.

USE DESIGNATED A ACRE UNLESS OTHER GROUND APPLICATI Crops — 60 to 800 g Use a minimum of

Where Recommended Crops — 10 to 20 g OBSERVE DAYS INT BY NUMBER IN () I

APPLES (30): EARLY European Red Mite, Dormant Oil Spray pe with Sulfur, Captan & SPRAYS — First Broc Leafminers—Use 1/4 day intervals in the per 100 gallons wa

BROCCOLI (3), CAU water to cover one control.

CANTALOUPE (1), W 1/4 to 1/2 pint per a COTTON (30): (Calif Aphids, Thrips, Fleah acre with ground eq Make first applicati

NET CONTENTS
GALLONS

SPRAY

BY WT.

.....	75.50%
.....	2.50%
.....	22.00%
.....	100.00%



REN

Skin! Wear a mask
phosphamidon pro-
skin or on clothing.
se of contact, wash
oroughly with soap
ing with soap and
ing areas or vicinity
be re-entered until
t contaminate feed

use inhibitor. Patient
idotal. Give 4 mg.
atient is fully atro-
(2-PAM) should be

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS WARNING AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS; WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE.

WARNING: Do not use on maple trees, peaches, plums and cherries as injury may occur. Avoid drift to these varieties.

Phosphamidon 8 Spray is a liquid systemic insecticide for the control of both sucking and chewing insects.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED.

GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gallons. **Deciduous Fruit Crops** — 60 to 800 gallons. **Citrus** — 400 to 3,000 gallons. (On young Fruit Plantings — Use a minimum of 15 gallons water per acre.)

Where Recommended for AIR APPLICATION — Field Crops — 3 to 10 gallons. **Orchard Crops** — 10 to 20 gallons.

OBSERVE DAYS INTERVALS BETWEEN LAST APPLICATION AND HARVEST INDICATED BY NUMBER IN () FOLLOWING CROP.

APPLES (30): EARLY DELAYED DORMANT ONLY — Green Peach and Rosy Apple Aphids, European Red Mite, San Jose Scale, Fruit Tree Leafroller — ¼ pint plus 1½ to 2 gallons Dormant Oil Spray per 100 gallons water as full coverage sprays. Do not combine oil sprays with Sulfur, Captan or Phaltan. **PRE-BLOSSOM AND POST-BLOSSOM FIRST BROOD COVER SPRAYS** — First Brood Codling Moth, Leafhoppers, Green Peach and Rosy Apple Aphids, Leafminers—Use ¼ pint per 100 gallons water. Repeat at 7 to 10-day intervals (14 to 21 day intervals in the West). Repeat application to prevent Aphid buildup — 1/6 pint per 100 gallons water.

BROCCOLI (3), CAULIFLOWER (3): Aphids, Imported Cabbageworm — 1 to 1 pint in water to cover one acre. Apply every 10-14 days in a preventative program for Aphid control.

CANTALOUPE (1), WATERMELON (1), CUCUMBERS (3): Aphids, Leafminer, Leafhopper — ¼ to ½ pint per acre in water to cover.

COTTON (30): (California and Arizona Only): **Early Season** (2 through 10 leaf stage) — Aphids, Thrips, Fleahoppers, Spider Mites — Apply 2 to 3 applications of 3 ounces per acre with ground equipment. Spray directly on young plants in sufficient water to cover. Make first application at the second true leaf stage. **Mid and Late Season** — Lygus,

Flea Beetles, Leafhoppers — 8 ounces per acre in water to cover, by ground or air. Aphids — 4 ounces per acre by ground or air. **Air Application** — Use 1 to 5 gallons water per acre. Do not apply more than 3 sprays. Do not graze treated cotton fields.

ORANGES (15), GRAPEFRUIT (15), LEMONS (15): Aphids, Thrips — ¼ to ½ pint per 100 gallons water (Ground Application — 1¼ to 2½ pints per acre in water to cover. By Air — ½ to ¾ pint per acre in 10 to 20 gallons water). Repeat as necessary. Apply during periods of Aphid or Thrips attack on new flush growth.

PEPPERS (6): Aphids, Leafminer, Whitefly — ½ pint per acre in water to cover. European Corn Borer — 1 pint per acre in water to cover. Apply when insects appear and when Corn Borer infestation is expected.

POTATOES (14): Aphids, Colorado Potato Beetles, Leafhoppers, Leafminer, Flea Beetles, Stinkbugs — ½ pint per acre in water to cover by ground application (1 to 5 gallons by air). Repeat as necessary. Tuberworm — ½ to 1 pint per acre in water to cover by ground (10 gallons by air). Apply prior to rows closing or at first sign of leaf mining. Repeat every 10 to 12 days.

TOMATOES (10): Aphids, Leafminer, Whitefly, Flea Beetles, Stinkbugs — ½ pint per acre in water to cover by ground application (1 to 5 gallons by air). Repeat as necessary.

WALNUTS (7): Aphids, Codling Moth — 1 pint in 400 gallons water per acre by ground application (10 to 20 gallons by air). Walnut Husk Fly — 1 pint in 200 to 300 gallons water per acre by ground (10 to 20 gallons by air). Apply after egg stings first occur on the husks.

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

