Code: 30062

LIQUI-STIK CONCENTRATE

Net Contents One Gallon

Liqui-Stik Concentrate is a hormone-like spray to be used in the following manner on apples and pears for the prevention of pre-harvest fruit drop and thinning of olives. Do not use in undiluted form.

DIRECTIONS FOR APPLES AND PEARS

Spray trees thoroughly, wetting all fruit stems and adjacent leaves. Apply 7 to 14 days before harvest for apples and 5 to 7 days before harvest for pears. Do not spray within 2 days of harvest. This product becomes effective in 2 to 3 days after application and controls drop for 10 days to 4 weeks (less for McIntosh variety) depending on weather conditions. Do not make more than 2 applications and do not delay picking beyond optimum maturity.

GROUND SPRAY: For Apples, thoroughly mix 12 fl. oz. with 250 gallons of water (4.8 fl. oz. per 100 gals, water) to prepare a 20 ppm. spray. 630 gallons per acre of this 20 ppm. spray provides an equivalent of 48 grams of 1-Naphthaleneacetic acid per acre. For Pears, use a 10 ppm. spray obtained by mixing 12 fl. oz. with 500 gallons of water, 670 gallons per acre of this 10 ppm. spray provides an equivalent of 25 grams of 1-Naphthaleneacetic acid per acre.

AIRPLANE SPRAY: For Apples USE 30 fl. on in 5 gallons of water. For Pears, use 11 ff. o water per acre.

025W

MANUFACTURED IN U.S.A. BY

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT

FMC CORPORATIONOR ECONOMIC POISON

AGRICULTURAL CHEMICAL DIVISION

MIDDLEPORT, N.Y. FRESNO, CA.

YAKIMA, WA RICHMOND, CA

| E.P.A. REGISTRATION NO. 279-2131 Z | Α |
|------------------------------------|---------|
| ACTIVE INGREDIENT: | BY WT. |
| 1-Naphthaleneacetic Acid as the | |
| ammonium salt* | 5.68% |
| INERT INGREDIENTS: | 94.32% |
| | 100.00% |

GROWTH REGULATOR

*Equivalent to 5.2% 1-Naphthaleneacetic acid or 200 grams per gallon.

CAUTION **KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Avoid contact of eyes and clothing by the diluted spray. Avoid contact of skin with the concentrate. Avoid contamination to feed and foodstuffs.

For Emergency Assistance Call: 716-735-3765

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply Where run-off is likely to occur. Apply this product only as specified on this label.

Do not re-use empty one gallon container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

DIRECTIONS FOR THINNING OLIVE TREES

Liqui-Stik Concentrate is effective in thinning olive trees when an excessively heavy fruit set is expected.

Timing is critical. Spray when young fruits are 1/8" to 3/16" in diameter, which is from 11/2 to 21/2 weeks following full bloom. Thinning is effective on all varieties except Sevillano. Without Oil. Apply a solution of 150 ppm. 1-Naphthaleneacetic acid equivalent obtained by adding 9 fl. oz. and a wetting agent (e.g. Dreft 3 to 4 oz.) to 25 gallons of water. With Oil. Apply a solution of 100 ppm. 1-Naphthaleneacetic acid equivalent obtained by mixing thoroughly 6 fl. oz. with 25 gallons water. Emulsify by agitation with 3 pints of light or medium summer oil. Spray only the heavily loaded trees until the leaves are thoroughly wet. This will require 10 to 30 gallons per tree, CAUTION: Unless you wish to remove all your fruit, do not apply during bloom.

Dealer Should Sell in Original Packages Only. WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use, EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS, FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.



E.P.A. EST., 279-CA-2

Printed in U.S.A.