

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of any body of water. Birds nesting in treated areas may be killed. Apply this product only as specified on this label. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is not to be used for the treatment of any animal. It is not to be used for the treatment of any plant or animal other than those listed on the label.

DIRECTIONS FOR USE

This material is a Federal law to use this product in a manner not specified on this label.

Apply the dosage indicated per acre if the proper amount of water is used for thorough coverage of the plant. Observe days interval between last application and harvest indicated by number in parentheses following crop.

Workers entering treated fields within 24 hours after application should wear protective clothing.

Peppers [15] Flea Beetle, Green Peach Aphid - Use 1 quart per acre.

Potatoes [5] Flea Beetle, Colorado Potato Beetle, Leafhopper, Aphids, Southern Armyworm, Green Stink Bug, Leaf-footed Plant Bugs and other Plant Bugs, Potato Tuberworm, Whitefly - Use 1 to 2 quarts per acre. False Chinch Bug - Use 2 quarts per acre. Potato Psyllid - Use 1 to 2 quarts per acre. Do not plant root crops other than carrots, potatoes, sugar beets, and sweet potatoes as follow-up crops.

Pumpkins [10] Aphids, Cucumber Beetles - Use 1 quart per acre. Do not apply before lining.

Tomatoes [10] Aphids, Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Whitefly - Use 1 quart per acre. Hornworms - Use 1.6 pints per acre.

Cotton Spider Mites (Desert Spider Mite, Two-spotted Spider Mite), Aphids, Cotton Leafworm - Use 1 quart per acre. Boll Weevil - Use 1.5 quarts per acre. Lygus Bug, Bollworm, Cabbage Looper - Use 2 quarts per acre. Do not apply after the bolls open. Do not graze dairy or meat animals in treated fields.

Tobacco [Field] [5] Tobacco Suckfly - Use 1 to 2 pints per acre. Budworm, Cabbage Looper, Flea Beetle, Green Peach Aphid, Hornworm - Use 1 quart per acre. Do not apply within 5 days of pruning or 15 days of cutting. Avoid contact with plants when pruning or cutting tobacco. Persons entering treated areas within 24 hours after application should wear protective clothing.

Tobacco [Seed Beds] In seed beds for control of the insects listed under Field Tobacco use 1 quart per 100 gallons of water. Apply 6 gallons of finished spray per 100 square yards of seed bed. For control of Green June Beetle Larvae in seed beds use 1 pint in 100 gallons of water. Apply 1 gallon of mixture per square yard of plant bed. Allow 15 days between last application and pulling plants.

STORAGE AND DISPOSAL

Rinse equipment and containers and DISPOSE OF WASTES by burying at least 18 inches deep in non-crop lands away from water supplies. Do not reuse containers, smash or perforate and bury with wastes.

Not for use or storage in or around the home

- (X) Do not contaminate feed or food stuffs.
 - (X) Do not use in or around homes.
 - (X) Do not store in homes.
 - (X) Do not use, pour, spill, or store near heat or open flame.
- Keep all unused material in original packages. Original packages should be removed from circulation.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: In purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather conditions, conditions mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are the responsibility of the seller and are assumed by buyer at the time of use.

Use of Product: FMC makes no representation for the use of this product in insect control tests, research, or otherwise. The use of this product being beyond the scope of the product label, no guarantee, expressed or implied, is made as to the results of such use. Results to be obtained if not used in accordance with the label instructions are the user's responsibility.

Damages: Buyer and user assume all liability for damages for breach of warranty, if required, shall be limited to direct damages not exceeding the purchase price paid for the product, and shall not include incidental or consequential damages.

