

Code: 30575

SULPHUR 50 THIODAN[®] 4 DUST

50 Pounds Net Weight

DIRECTIONS FOR USE

Unless indicated otherwise apply when pests or disease first appear AND REPEAT AS NECESSARY. Application may be made with ground or airplane application equipment.

ON APPLES to control Apple Aphid and Powdery Mildew use 45-80 pounds per acre. At the 45-60 pounds per acre rate with no limitation on the number of applications, an application may be made within 30 days of harvest. At the 45-60 pounds per acre rate if no more than 3 applications are made during the fruiting period an application may be made within 21 days of harvest. At a rate greater than 60 pounds do not make more than 2 applications nor apply within 30 days of harvest. Sulfur may injure certain varieties, check tolerance prior to use. DO NOT FEED POMACE FROM TREATED APPLES TO LIVE-STOCK.

ON ARTICHOKE to control Artichoke Aphid, Green Peach Aphid and Artichoke Tume Moth use 25-35 pounds per acre. At the 25 pound per acre rate do not apply within 7 days of harvest. At a rate exceeding 25 pounds do not apply after buds start to form.

ON BEANS to control Bean (Black Bean) Aphid, Bean Leaf Skeletonizer, Cucumber Beetles, Green Stink Bug and Mexican Bean Beetle use 25 pounds per acre. Do not apply more than 3 times per season. Do not apply within 3 days of harvest. Do not feed treated threshings to livestock or allow to graze in treated fields. Do not use on lima beans that are to be removed from the field for processing. Sulfur may injure certain varieties, check sulfur tolerance prior to use.

ON BROCCOLI, CABBAGE, BRUSSELS SPROUTS and CAULIFLOWER to control Cabbage Aphid, Cabbage Looper, Cross-striped Cabbageworm, Diamond-back moth larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm and Stink Bug use 25 pounds per acre. On Broccoli and Cabbage do not apply within 7 days of harvest. On Brussels Sprouts and Cauliflower do not apply within 14 days of harvest.

ON CARROTS to control Green Peach Aphid use 25 pounds per acre. Do not make more than 2 applications nor apply within 7 days of harvest. Do not use tops for food or feed.

ON CELERY to control Green Peach Aphid and Cabbage Looper use 25 pounds per acre. Do not apply more than 3 times per season. Do not apply within 4 days of harvest.

ON GRAPES to control Grape Leafhopper, Powdery Mildew and Grape Leaf Folder (Calif. Only) use 25-35 pounds per acre. Do not apply within 7 days of harvest. Do not apply to Concord variety as severe injury is likely to occur. Ground application equipment is preferred.

ON HEAD LETTUCE to control Cabbage Looper, Diamondback moth larvae, Green Peach Aphid and Imported Cabbageworm use 25 pounds per acre. Do not apply more than 3 times after thinning. Do not apply within 14 days of harvest. Remove outside leaves at harvest. Do not feed crop waste to livestock.

ON PEACHES and NECTARINES to control Brown Rot, Powdery Mildew, Green Peach Aphid and Peach Twig Borer use 40-80 pounds per acre. Do not make more than 2 applications during the fruiting period. Do not apply within 30 days of harvest.

ON PEARS to control Apple Aphid and Apple Rust Mite use 40-60 pounds per acre. Do not apply more than twice during fruiting season. Do not apply within 7 days of harvest. Sulfur may injure certain varieties, check tolerance prior to use.

FUNGICIDE - INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-2213

ACTIVE INGREDIENTS:	BY WT.
*Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide) - - - - -	4.00
Sulphur - - - - -	50.00%
INERT INGREDIENTS - - - - -	46.00%
	100.00%

* Thiodan is a registered trademark of Canadian Hoechst Ltd. - U.S. Patent No. 2,799,685

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas on the day of application should wear protective clothing.

USAGE CAUTION

Applications of materials containing sulphur should not be made prior to or during periods of excessively high temperatures.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Apply this product only as specified on the label.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not allow this material to drift onto sulfur sensitive crops.

ON GARDEN PEAS (Seed only) to control Pea Aphid and Pea Weevil use 25 pounds per acre. Do not feed treated vines or threshings to meat or dairy animals.


ON PLUM and PRUNES to control Peach Twig Borer and Brown Pot use 50-60 pounds per acre. Do not apply within 7 days of harvest.

ON STRAWBERRIES to control Meadow Spittlebug, Strawberry Aphid and Powdery Mildew use 25 pounds per acre. Do not reapply within 15 days or more than twice within a 35 day period when fruit is present. Do not apply within 4 days of harvest.

ON SWEET CORN (FOR FRESH VEGETABLE USE ONLY) to control Corn Leaf Aphid use 25-35 pounds per acre. To control Corn Earworm use 35 pounds per acre. For Corn Earworm apply at 2 to 3 day intervals beginning when silks first appear and continuing until they begin to dry. Do not apply more than 5 times. Do not apply to sweet corn to be processed. Do not feed treated forage or animals to livestock or allow to graze in treated fields.

ON TOMATOES to control Blister Beetles, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Cabbage Looper, Stink Bug, Tomato Fruitworm, Yellow-striped Armyworm, Tomato Hornworm, Tomato Russet Mite and Whitefly use 25 pounds per acre. Do not apply within 1 day of harvest.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives all claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS In the event of certain critical conditions created by conditions beyond the control of Niagara, which Niagara has no control over, as a condition of purchase, user assumes responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such cause. No person is authorized to vary or negate any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

