

Code 2603



Net Contents
Gallons

Ethion 1.0 Tedion 0.75 EC

Miticide

EPA Reg. No. 279-2226

EPA Est., 279-

Active Ingredient:	By Wt.
Tetradifon (4-chlorophenyl 2,4,5-trichlorophenyl sulfone)	8.8%
*Ethion	11.6%
Xylene base aromatic petroleum distillate ...	31.0%
Inert Ingredients:	48.6%
	100.0%


*O,O,O'-Tetraethyl S,S'-methylene bisphosphorodithioate
Contains 1.0 lbs Ethion and 0.75 lb Tedion per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

For Emergency Assistance Call 716-735-3765.

See Other Panels for Additional Precautionary Information.



FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia, Pennsylvania 19103

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution
Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products.

Environmental Hazards

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Physical or Chemical Hazards

Flammable. Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not store below 32°F. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Keep container closed. Leaking packages should be removed to a safe place. Do not reuse container. To dispose of container, triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. Do not cut or weld this container. Consult Federal, State or local disposal authorities for approved alternative procedures.

Do not use in undiluted form.

Thoroughly mix in sufficient water per 100 gallons or amount required to treat one acre. When using this material in the same spray tank with wettable powders, add the wettable powders first and after they are thoroughly dispersed in the tank, add this material to the tank last. Apply when mites first appear and repeat as required to maintain control. Apply to both upper and lower leaf surfaces. Observe use limitations. Observe days interval between last application and harvest indicated by number in () following crop.

Apple (40): European Red Mite, Twospotted Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite, Brown Almond Mite—Use 1 qt. per 100 gals. Do not apply more than 4 applications during the fruiting period, and do not apply more than 500 gals. of spray per acre per application.

Apple (20): European Red Mite, Twospotted Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite, Brown Almond Mite—Use 1 qt. per 100 gals. Do not apply more than 2 applications during the fruiting period, and do not apply more than 500 gals. of spray per acre per application.

Do not use on Astrachen, Davey, Duchess, Margaret Praline, Wealthy, Williams Red (Red Williams), Yellow Transparent varieties maturing at or before early McIntosh or on Crimson #3 (Brite Mac) apples as injury is likely to occur.

Do not feed pomace from treated apples to livestock.

Cherry: European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite—Use 1 qt. per 100 gals. Do not use more than 2 applications during the fruiting period. Use as a post-harvest treatment only.

Ornamentals (Out-of-Doors): Twospotted Mite—Use 1 qt. per 100 gals. of water. Citrus Red Mite—Use 1 1/2 qts. per 100 gals. of water. Do not apply to Anderson Yew. Application of this material to Butterfly and Cinderella roses may cause leaf drop. Do not open Chrysanthemum blossoms (any variety) as slight injury may occur.

Peach (30): European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite—Use 1 qt. per 100 gals. Do not apply more than 2 applications during the fruiting period. Do not apply more than 500 gals. of spray per acre per application.

Pear (30): European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite, Canadensis Mite, Carpini Mite, Schoenii Mite—Use 1 qt. per 100 gals. Do not apply more than 500 gals. of spray per acre per application. Do not apply more than 3 applications, first cover spray.

Plums (Fresh Prunes) (21): European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite—Use 1 qt. per 100 gals. Do not make more than 3 applications during the fruiting period. Do not apply more than 600 gals. of spray per acre per application.

Dealers Should Sell in Original Packages

Terms of Sale or Use: On purchase of this product buyer agrees to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, in the use of this product other than indicated on the label. Except as warranted, the product is sold as is. Buyer and user assume all use and/or handling and/or storage of this material where use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully and method of application, weather and crop conditions, and other chemicals not specifically recommended and other use factors in the use of this product are beyond the control of FMC and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or other safe practice.

Damages: Buyer's or user's exclusive remedy for damages, in whole or in part, of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include consequential damages.



Net Contents Gallons

on 1.0 Tedion 0.75 EC

1-2223 EPA Est., 279-

	By Wt.
chlorophenyl 2,4,5-trichlorophenyl	8.8%
.....	11.6%
omatic petroleum distillate ...	31.0%
.....	48.6%
	<u>100.0%</u>

S-methylene bisphosphorodithioate
and 0.75 lb Tedion per gallon

OF REACH OF CHILDREN

CAUTION

ince Call 716-735-3765.

Additional Precautionary Information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution
Harmful if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products.

Environmental Hazards

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Physical or Chemical Hazards

Flammable. Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not store below 32°F. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Keep container closed. Leaking packages should be removed to a safe place. Do not reuse container. To dispose of container, triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place. Do not cut or weld this container. Consult Federal, State or local disposal authorities for approved alternative procedures.

Do not use in undiluted form.

Thoroughly mix in sufficient water per 100 gallons or amount required to treat one acre. When using this material in the same spray tank with wettable powders, add the wettable powders first and after they are thoroughly dispersed in the water, add this material to the tank last. Apply when mites first appear and repeat as required to maintain control. Apply to both upper and lower leaf surfaces. Observe use limitations. Observe days interval between last application and harvest indicated by number in () following crop.

Apple (40): European Red Mite, Twospotted Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite, Brown Almond Mite—Use 1 qt per 100 gals. Do not apply more than 4 applications during the fruiting period, and do not apply more than 500 gals of spray per acre per application.

Apple (20): European Red Mite, Twospotted Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite, Brown Almond Mite—Use 1 qt per 100 gals. Do not apply more than 2 applications during the fruiting period, and do not apply more than 500 gals of spray per acre per application.

Do not use on Astrachen, Davey, Duchess, Margaret Pratt, Melba, Wealthy, Williams Red (Red Williams), Yellow Transparent or any varieties maturing at or before early McIntosh or on Crimson and N J #3 (Brite Mac) apples as injury is likely to occur.

Do not feed pomace from treated apples to livestock.

Cherry: European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite—Use 1 qt per 100 gals. Do not use more than 500 gals of spray per acre per application. Use as a post-harvest treatment only.

Ornamentals (Out-of-Doors): Twospotted Mite—Use 1 qt per 100 gals of water. Citrus Red Mite—Use 1 1/2 qts per 100 gals of water. Do not apply to Anderson Yew. Application of this material to White Butterfly and Cinderella roses may cause leaf drop. Do not apply to open Chrysanthemum blossoms (any variety) as slight injury may occur.

Peach (30): European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite—Use 1 qt per 100 gals. Do not apply more than 2 applications during the fruiting period. Do not apply more than 500 gals of spray per acre per application.

Pear (30): European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite, Canadensis Mite, Carpini Mite, Schoenii Mite—Use 1 qt per 100 gals. Do not apply more than 500 gals of spray per acre per application. Do not apply more than 3 applications after the first cover spray.

Plums (Fresh Prunes) (21): European Red Mite, Twospotted Mite, McDaniel Mite, Brown Almond Mite—Use 1 qt per 100 gals. Do not make more than 3 applications during the fruiting period. Do not apply more than 600 gals of spray per acre per application.

Dealers Should Sell in Original Packages Only.


Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and or handling and or storage of this material when such use and or handling and or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.



tion
Chemical Group
Street
Pennsylvania 19103