

Code: 31643



Gallons N

PARATHION 8 F

DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary. Use the following recommended rates in sufficient water to thoroughly cover one acre. (By ground, apply a minimum of 200 gallons per acre on stone fruits and nuts; a minimum of 300 gallons per acre on pome fruits; a minimum of 200 gallons per acre on bush and vine fruits and a minimum of 10 gallons per acre on vegetable and row crops. By air, apply a minimum of 2 gallons per acre on vegetable and field crops and a minimum of 20 gallons per acre on small fruits and tree crops.)

ON APPLES and PEARS to control Codling Moth, Orange Tortrix, San Jose, Parlatoria and Forbes Scales, Woolly Apher, Mealy Bug, Bud Moth, Fruit Tree Leafroller, Leafhoppers, Rosy-Apple Aphid and Green Apple Aphid use $\frac{1}{4}$ - $\frac{1}{3}$ pint per 100 gallons. By airplane, $\frac{1}{2}$ pints per acre. **ON PEAR** to control Blister Mite and Pear Psylla use $\frac{1}{4}$ - $\frac{1}{3}$ pint per 100 gallons in pre-blossom and blossom sprays. By airplane, $\frac{1}{2}$ - $\frac{1}{3}$ pints per acre. May cause injury to fruit and foliage of McIntosh and related varieties, to Golden Delicious and to Bosc Pears. Do not apply within 14 days of harvest.

ON CHERRY to control Cherry Fruitworm, Oriental Fruit Moth, Bud Moth, Orange Tortrix, Cankerworm, Fruit Fly, San Jose Scale Crawlers and Aphids use a pint per 100 gallons. By airplane, 1 pint per acre. Do not apply within 14 days of harvest.

ON FIG to control Fig Scale use a pint per 100 gals. Do not apply within 30 days of harvest.

ON PLUMS and PRUMES to control Peach Twig Borer, Orange Tortrix, Leafhopper, Codling Moth, Tussock Moth, Bud Moth, Mealy Plum Aphid, Thrips, Fruit Tree Leafroller and San Jose Scale use $\frac{1}{4}$ - $\frac{1}{3}$ pint per 100 gallons. By airplane $\frac{1}{2}$ - $\frac{1}{3}$ pints per acre. Do not apply within 14 days of harvest.

ON NECTARINES and PEACHES to control Oriental Fruit Moth, Peach Twig Borer, Aphid, Fruit Tree Leafroller and Can Facing Insects use a pint per 100 gallons (Apply a minimum of 1 quart per acre). To control Thrips use $\frac{1}{4}$ - $\frac{1}{2}$ pint per 100 gals. Apply a min. of 1 quart per acre. **CALIFORNIA:** Do not apply more than 5 pounds of actual Parathion per acre between January 1 and harvest. Do not apply more than once after bloom. Do not apply within 21 days of harvest. **AREAS OTHER THAN CALIFORNIA:** Do not apply more than 5 pounds of actual Parathion per acre per year. Do not apply within 14 days of harvest.

ON GRAPES to control Thrips, Hopper, Beetle, Mealy Bug, Leafroller use $\frac{1}{4}$ - $\frac{1}{3}$ pint per 100 gallons (200 gallons per acre). Do not apply within 14 days of harvest. Grape Mealy Bug: Dormant application use 1x pint plus 2 gallons Dormant Oil Emulsion per 100 gallons. Apply 200 gallon per acre as a dormant spray. Apply only before fruit is size of buckshot or after harvest. Apply with ground application equipment only.

ON BEANS to control Corn Earworm, Diabrotica Beetles, Leafhoppers, Aphids, Leafminers, Whitefly and Armyworm use $\frac{1}{4}$ - $\frac{1}{3}$ pint per acre. Do not apply within 14 days of harvest.

INSECTICIDE

U.S.D.A. REGISTRATION 279-2230

ACTIVE INGREDIENT:
Parathion
INERT INGREDIENTS:

BY WT.
80.00%
20.00%
100.00%

0,0-Diethyl 0-p-nitrophenyl thiophosphate
Contains 8.0 Pounds Parathion Per Gallon



DANGER-POISON



KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

FIRST AID TREATMENT

ANTIDOTE: If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY.** If on skin, in case of contact, remove contaminated clothing and immediately wash skin with soap and water.

Parathion is a cholinesterase inhibitor and treatment of poisoning should include the use of atropine.

Keep on hand an emergency supply of 20 tablets, grains 1/100. In case of poisoning take 3 tablets, grains (1/100) and call a physician at once.

PHYSICIAN: Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

BEFORE USING SHAKE OR ROLL CONTAINER.

Do not store below 0 F.

FLAMMABLE. KEEP AWAY FROM FIRE, HEAT AND OPEN FLAME. LEAKING PACKAGES SHOULD BE REMOVED TO A SAFE PLACE.

DO NOT USE IN UNDILUTED FORM

Manufactured in U.S.A. by

NIAGARA CHEMICAL DIVISION
FMC CORPORATION

MIDDLEBURY, N. Y.



FRESNO, CALIF. • RICHMOND, CALIF.
RIVERSIDE, CALIF. • YAKIMA, WASHINGTON

ON STRAWBERRIES to control Aphids, berry Leafroller, Root Weevil and 1 gallon. By airplane, $\frac{1}{2}$ pint per acre of harvest.

ON LETTUCE to control Aphids, Leaf Armyworms use $\frac{1}{2}$ pint per acre. On 7 days; On Leaf Lettuce do not apply. **ON SPINACH** to control Aphids, Leaf and Seed Corn Maggot in Crowns use within 14 days of harvest.

WARNING: Poisonous if swallowed, skin! Rapidly absorbed through skin or on clothing. Wear natural rubber and goggles. In case of contact, wash with water. Wear a mask or respirator. Department of Agriculture for unprotected persons and beneficial or vicinity where there may be bees should not be re-entered until dry residues have dissipated. Do not eat. Wash hands, arms and face thoroughly before eating or smoking. Wash all contact with hot water before re-use.

USAGE CAUTION

Do not apply when bees are active. This product is poisonous to fish, lakes or ponds. Do not apply in water.

Rinse equipment and container and least 16 inches deep in non crop. Empty containers should be smashed or burned.

Do not combine with emulsifiable. proven them to be compatible.

Dealer Should Sell in Original

USAGE CAUTION: Do not allow this material to come in contact with or use in a manner or at a time other than that specified on the label. animal, plant or crop injury, illegal residues or other. **TERMS OF SALE OR USE:** On purchase or use of this product, the following conditions apply.

WARRANTY: Niagara warrants that this product, conforms to the ingredient statement on the label, and in the Directions for Use. EXCEPT AS SO WARRANTED, NIAGARA MAKES NO OTHER WARRANTY. **DAMAGES:** Buyer's or user's exclusive remedy for damages shall be limited to direct damages not exceeding incidental or consequential damages. Notice within 10 days of discovery by buyer or user, for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: created or incurred by buyer or user or over which as a condition of purchase or use, assume responsibility arising out of the handling or use of this person is authorized to vary or waive any state.



Gallons Net Contents

PARATHION 8 F

INSECTICIDE

U.S.D.A. REGISTRATION 279-2230

ACTIVE INGREDIENT :
Parathion
INERT INGREDIENTS:

BY WT.
80.00%
20.00%
100.00%

O,O-Diethyl O-p-nitrophenyl thiophosphate
Contains 8.0 Pounds Parathion Per Gallon



DANGER-POISON



KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

FIRST AID TREATMENT

ANTIDOTE. If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY.** If on skin, in case of contact, remove contaminated clothing and immediately wash skin with soap and water.

Parathion is a cholinesterase inhibitor and treatment of poisoning should include the use of atropine.

Keep on hand an emergency supply of 20 tablets, grains 1/100. In case of poisoning take 3 tablets, grains (1/100) and call a physician at once.

PHYSICIAN: Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

BEFORE USING SHAKE OR ROLL CONTAINER.

Do not store below 0 F.

FLAMMABLE: EXCEPT BY THE HEAT AND OPEN FLAME. LEAKING PACKAGES
DO NOT USE IN UNDILUTED FORM

DO NOT USE IN UNDILUTED FORM

Manufactured in U.S.A. by

NIAGARA CHEMICAL DIVISION

FMC CORPORATION

MIDDLEBURY, N. Y.



FRESNO, CALIF.

RICHMOND, CALIF.

DIVERSIDE, CALIF.

YAKIMA, WASHINGTON



ON STRAWBERRIES to control Aphids, Lygus Bugs, Leafhoppers, Strawberry Leafroller, Root Weevil and Whitefly use 1/4-1/3 pint per 100 gallons. By airplane, 1/2 pint per acre. Do not apply within 14 days of harvest.

ON LETTUCE to control Aphids, Leafhoppers, Leafminer, Loopers and Armyworms use 1/2 pint per acre. On Head Lettuce do not apply within 7 days; On Leaf Lettuce do not apply within 21 days of harvest.

ON SPINACH to control Aphids, Leafminer, Crown Mite (Calif.) and Seed Corn Maggot in Crowns use 1/2 pint per acre. Do not apply within 14 days of harvest.

WARNING: Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for Parathion protection. Keep all unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

USAGE CAUTION

Do not apply when bees are active in the field. This product is poisonous to fish and wildlife. Keep out of streams, lakes or ponds. Do not apply in any manner not specified on this label.

Rinse equipment and container and dispose of wastes by burying at least 16 inches deep in non crop land away from water supplies. Empty containers should be smashed and buried with wastes. Do not burn.

Do not combine with emulsifiable liquids unless previous use has proven them to be compatible.

Dealer Should Sell in Original Packages Only.

USAGE CAUTION Do not allow this material to drift onto neighboring crops or noncrop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

TERMS OF SALE OR USE On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purpose set forth in the Directions for Use EXCEPT AS SO WARRANTED. THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user as a condition of purchase or use assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.