

Code: 30412

SEVIN* SULPHUR 10-50 DUST

DIRECTIONS FOR USE

Apply when insects or disease first appear and repeat as necessary for insect or disease control beyond the time limitations of this product. Use a suitable material. May be applied with ground or airplane equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

ALMONDS (0): Codling Moth, Filbert Worm, Fruit Tree Leafroller, Brown Rot. Use 10 to 80 pounds per acre.

APPLES (1), PEARS (1): Powdery Mildew, Codling Moth, Green Apple Aphid, Apple Maggot, Red-Banded Leafroller, Pear Psylla, Periodical Cicada—Use 40 to 80 pounds per acre. Application within 30 days after full bloom may also provide apple thinning. To avoid this, delay use until at least 30 days after bloom.

ASPARAGUS (1): Asparagus Beetle on Seedlings or Spears. Use 20 pounds per acre. Apply during cutting season. Do not repeat application within 3 days. Asparagus Beetle, Apache Cicada on ferns or brush growth in the post harvest period. Use 20 to 40 pounds per acre.

BEANS (0): Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Leafhoppers. Use 10 pounds per acre. Corn Earworm, Stink Bugs, Tarnished Plant Bug. Use 15 pounds per acre. California Lygus Bug, Stink Bug, Lima Bean Pod Borer. Use 20 pounds per acre. Sulfur may cause injury to certain bean varieties, check sulfur tolerance prior to using.

BLACKBERRIES (7), BOYSENBERRIES (7), DEWBERRIES (7), LOGANBERRIES (7), RASPBERRIES (7): Leafrollers, including Omnivorous Leafrollers, Sawfly, Climbing Cutworms, Snowy Tree Cricket (Calif.), Orange Tortrix (Oregon), Leafhoppers, Raspberry Aphid, Powdery Mildew. Use 20 pounds per acre.

CABBAGE (3), BROCCOLI (3), BRUSSELS SPROUTS (3), CAULIFLOWER (3), KOHLRABI (3): Flea Beetles, Harlequin Bugs. Use 10 pounds per acre. Armyworm, Corn Earworm, Cabbage Looper (small). Use 20 pounds per acre. For small Cabbage Looper control after edible parts start to form, apply in a 5 to 7 day application.

CARROTS (0), CHINESE CABBAGE (14), COLLARDS (14), DANDELION (14), ENDIVE (14), BEETS (Roots 3) (Tops 14), HORSERADISH (3), KALE (14), LETTUCE (Head 3) (Leaf 14), MUSTARD GREENS (14), PARSLEY (14), PARSNIPS (3), RADISHES (3), RUTABAGAS (3), SALSIFY (Roots 3) (Tops 14), SPINACH (14), SWISS CHARD (14), TURNIPS (Roots 3) (Tops 14): Flea Beetle, Harlequin Bugs, Leafhoppers. Use 10 pounds per acre. Six-Spotted Leafhopper. Use 15 pounds per acre. Armyworm, Corn Earworm, Tarnished Plant Bug, Stink Bugs, Cabbage Looper (small). Use 20 pounds per acre. For small Cabbage Looper control apply in a 5 to 7 day application schedule.

CHERRIES (1), PLUMS (1), PRUNES (1): Cherry Fruit Fly, Peach Twig Borer, Leafhopper, Codling Moth, Orange Tortrix, Western Tussock Moth, Brown Rot, Powdery Mildew. Use 40 to 60 pounds per acre.

CORN (0): Corn Earworm, Fall Armyworm, Flea Beetles, Sap Beetles, Leafhoppers. Use 20 pounds per acre. For larvae in whorl and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when silks appear and continuing until silks begin to dry. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

073W

FUNGICIDE-INSECTICIDE

E.P.A. REGISTRATION NO. 279-2260 AA

ACTIVE INGREDIENTS	BY WT.
Carbaryl, (1-Naphthyl N-methylcarbamate)	10.00%
Sulphur	50.00%
INERT INGREDIENTS	40.00%
	100.00%

*Trade-mark Union Carbide Corporation for the active ingredient carbaryl.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

USAGE CAUTIONS

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use with Lime, Bordeaux, Lime Sulphur, Casein-Lime, Spreaders or other alkaline materials.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

Applications of materials containing sulphur should not be made prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Do not apply when weather conditions favor drift from the areas treated. Do not apply when run off is likely to occur.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765

COTTON (0): Early Season: Thrips, Flea B, Striped Blister Beetle. Use 10 pounds per acre. Bollworm, Fall Armyworm, Cotton Leaf P, Plant Bug, Lygus Bug, Aphids (suppression worm). Use 20 to 25 pounds per acre. Stink bugs per acre. Grasshoppers. Use 10

FILBERTS (0): Filbert Aphid, Filbert Leafhopper.

GRAPES (0): Powdery Mildew, Grape Leafhopper. Use 20 pounds per acre. Apply after emerge from buds and as needed for Leaf

PEACHES (1), NECTARINES (3): Oriental Fruit Moth, Codling Moth, Tussock Moth. Use 40 to

OKRA (0): Corn Earworm, Stink Bugs. Use 10

RICE (14): Leafhoppers, Armyworms, Tadpole. Do not apply within 14 days before harvest. (Stam F 34 or Rogue) as injury may result.

STRAWBERRIES (1): Powdery Mildew, Strawberry Weevil. Use 20 pounds per acre.

SUGAR BEETS (14): Armyworms, Leafhopper, Grasshoppers. Use 15 pounds per acre.

TOMATO (0), EGGPLANT (0), PEPPERS (0): Leafhoppers. Use 10 pounds per acre. Fall Armyworm, Tarnished Plant Bugs, Russet M, etc.

VIN* SULPHUR 10-50 DUST

Net Weight
50 Pounds

FUNGICIDE-INSECTICIDE

E.P.A. REGISTRATION NO. 279-2260 AA

ACTIVE INGREDIENTS:	BY WT.
Carbaryl (1-Naphthyl N-methylcarbamate)	10.00%
Sulphur	50.00%
INERT INGREDIENTS:	
	40.00%
	100.00%

*Trademark Union Carbide Corporation for the active ingredient Carbaryl.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

USAGE CAUTIONS

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use with Lime, Bordeaux, Lime Sulphur, Casein-Lime Spreaders or other alkaline materials.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

Applications of materials containing sulphur should not be made prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765

COTTON (0): Early Season: Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Striped Blister Beetle. Use 10 pounds per acre. **Mid and Late Season:** Boll Weevil, Bollworm, Fall Armyworm, Cotton Leaf Perforator, Leafrollers, Leafhoppers, Tarnished Plant Bug, Lygus Bug, Aphids (suppression). Use 10 to 20 pounds per acre. Pink Bollworm. Use 20 to 25 pounds per acre. Stink Bug, Salt Marsh Caterpillar. Use 20 to 25 pounds per acre. Grasshoppers. Use 10 to 15 pounds per acre.

FILBERTS (0): Filbert Aphid, Filbert Leafroller, Filbert Moth. Use 40 to 50 pounds per acre.

GRAPES (0): Powdery Mildew, Grape Leafroller, Grape Leafhopper, Grape Leaf Skeletonizer. Use 20 pounds per acre. Apply just before first brood Leafroller larvae emerge from rolls and as needed for Leafhoppers. Ground application preferred.

PEACHES (1), NECTARINES (3): Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth. Use 40 to 60 pounds per acre.


OKRA (0): Corn Earworm, Stink Bugs. Use 20 pounds per acre.

RICE (14): Leafhoppers, Armyworms, Tadpole Shrimp (California). Use 15 pounds per acre. Do not apply within 14 days before or after applications of the herbicide propanil (Stam F-34 or Rogue) as injury may result.

STRAWBERRIES (1): Powdery Mildew, Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil. Use 20 pounds per acre.

SUGAR BEETS (14): Armyworms, Leafhoppers, Webworms. Use 20 pounds per acre. Grasshoppers. Use 15 pounds per acre.

TOMATO (0), EGGPLANT (0), PEPPERS (0): Colorado Potato Beetle, Flea Beetle, Leafhoppers. Use 10 pounds per acre. Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bugs, Russet Mites, Stink Bugs. Use 20 to 40 pounds per acre.



Printed in U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.

LABEL

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals