

DIRECTIONS FOR USE
Apply when pests or disease first appear and repeat as necessary. For control beyond the usage limitations of this product use a suitable material. May be applied with airplane or ground application equipment.

ON COTTON to control Aphids, Boll Weevil, Brown Cotton Leafworm, Cotton Fleahopper, Cotton Leafworm, Lygus Bugs, Mites, Thrips, Rapid Plant Bug, Tarnished Plant Bug, Stink Bug, Bollworm and Pink Bollworm use 16 to 33 pounds per acre. Do not apply within 1 day of picking. If late season applications are made, do not graze livestock on treated areas or feed gin wastes.

ON APPLES, CRABAPPLES, PEARS AND QUINCES to control Aphids, Apple Maggot, Codling Moth, European Apple Sawfly, Eye Spotted Bud Moth, Fruit Tree Leaf Folder, Leafhoppers, Mites, Red banded Leafroller and Powdery Mildew use 35-70 pounds per acre. Do not apply more than 8 times per season nor within 15 days of harvest.

ON CHERRIES to control Eye-Spotted Bud Moth, Fruit Tree Leafroller, Mites, Plum Curculio and Brown Rot use 35-80 pounds per acre. Do not apply more than 8 times per season nor apply within 15 days of harvest.

ON NECTARINES AND PEACHES to control Aphids, Mites, Platynota Flavescens Leafroller, Red-banded Leafroller, Peach Twig Borer (California Only) Powdery Mildew and Brown Rot use 35-80 pounds per acre. Do not apply more than 8 times per season nor within 21 days of harvest.

ON PLUMS AND PRUNES to control Aphids, Eye-spotted Bud Moth, Fruit Tree Leafroller, Mites, Red-banded Leafroller, Peach Twig Borer (California Only) and Brown Rot use 35-80 pounds per acre. Do not apply more than 8 times per season nor apply within 15 days of harvest.

ON BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES and RASPBERRIES to control Aphids, Leafhoppers, Leafminers, Leafrollers, Obscure root weevil and Powdery Mildew apply 20-30 pounds per acre. Do not apply within 14 days of harvest.

ON GRAPES to control Grape Berry Moth, Leafhoppers, Mites, Thrips (California) and Powdery Mildew use 20-30 pounds per acre. To control Red banded Leafroller use 25-30 pounds per acre. Do not apply more than 3 applications per season nor within 28 days of harvest.

ON BEANS to control Aphids, Bean Leafbeetle, Leafhoppers, Leafminers, Leafrollers, Mites and Stink Bugs use 15-20 pounds per acre. To control Mexican Bean Beetle, Spotted Cucumber Beetle, Striped Cucumber Beetle, Tarnished Plant Bug, Western-striped Cucumber Beetle and Powdery Mildew use 20 pounds per acre. Do not apply to Snap Beans within 7 days of harvest. Do not apply to dry Beans within 30 days of harvest, nor apply more than 4 applications. Do not feed ensile or treated forage. Sulphur may injure certain bean varieties. Check tolerance to sulphur prior to use.

ON BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND CAULIFLOWER to control Aphids, Cabbage Looper, Diamondback Moth, and Imported Cabbageworm use 20-25 pounds per acre. Do not apply within 7 days of harvest for Brussels Sprouts, 15 days of harvest for broccoli and cauliflower, nor within 21 days of harvest for cabbage.

ON COWPEAS (Southern Peas, Blackeyed peas, Crowder Peas) to control Leafminers and Stink Bugs use 15-33 pounds per acre. To control Corn Earworm and Cowpea Curculio use 33 pounds per acre. Do not apply more than 4 times per season nor within 7 days of harvest. Do not use vines for feed or forage nor pasture treated areas. Sulphur may injure certain varieties, check tolerance prior to use.

ON POTATOES to control Colorado Potato Beetle and Leafminers use 15-25 pounds per acre. To control Tuberworm, Aphids, European Cornborer, Flea Beetles, Leafhoppers, Scittlebugs and Tarnished plant bug use 20-25 pounds per acre. Do not apply within 7 days of harvest.

ON SPINACH to control Aphids, Leafminers and Mites use 15-25 pounds per acre. Do not apply within 14 days of harvest.

CODE
30232

50 POUNDS
NET WEIGHT

KOLODUST® 50 G-3 DUST

CONTAINS 95% GUTHION
INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-2261

ACTIVE INGREDIENTS:

BY WT.

0,0-Dimethyl S-(4-oxo-1,2,3-benzotriazin-3 (4H) ylmethyl) phosphorodithioate - 3.00%
Sulphur - 42.00%

INERT INGREDIENTS - 55.00%

† U.S. Patent No. 2,758,115
Canadian Patent No. 552,660

100.00%



DANGER - POISON



KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

USAGE CAUTION

Do not graze livestock in treated areas for 21 days after treatment.

Do not treat food crops grown in the green house. Dispose of wastes by burying in non-crop land away from water supplies.

Empty containers should be disposed of by burying with wastes or by burning. Keep Out of Smoke.

Highly toxic to fish. Do not contaminate any body of water nor apply to any area not specified on the label. Applications of materials containing Sulphur should not be made just prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Guthion is a registered trademark, Reg. U.S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G., Chemagro Corporation Licensee.

REVISED

ON TOMATOES to control Colorado Potato Beetle, Green S. Drosophila and Leafminers use 15-25 pounds per acre. To Aphids, European Corn borer, Flea Beetles, Hornworm, Leafhopper, Thrips use 20-25 pounds per acre. To control Corn Earworm, Pinworm, Tuberworm, and Yellow-striped Armyworm use 15-25 pounds per acre. No time limitations.

ON DRY ONIONS to control Thrips use 20-25 pounds per acre apply within 28 days of harvest. Do not apply more than 3 times per season.

ON ORNAMENTALS and NURSERY PLANTS to control Aphids, Lacebugs, Hoppers, Mites and Thrips use 20 pounds per acre. Dust all surfaces including the underside of leaves for complete coverage. Do not allow children or pets on treated areas until material washed into soil and treated area is dry.

WARNING

Poisonous if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Wear protective clothing, rubber gloves, and goggles.

Do not breathe dust. Wear a mask or respirator effective protection against GUTHION. Information as to suitable masks or respirators is available from the U.S. Department of Agriculture.

Do not contaminate feed or food. Keep out of reach of children. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Workers who use treated fields within 2 days of application should wear protective clothing. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest, sweating, contracted pupils, stomach pains, vomiting, diarrhea.

TREATMENT: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomit by giving a tablespoonful of salt in a glass of water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

ACCEPTED

MAR 13 1968

U.S. DEPARTMENT OF AGRICULTURE
FEDERAL AGENCY FOR PEST CONTROL
WASHINGTON, D.C. 20250
REGISTRATION NO. 279-2261

Printed

CODE
30232

50 POUNDS
WET WEIGHT

KOLODUST® 50 G-3 DUST

**CONTAINS 3% GUTHION
INSECTICIDE**

U.S.D.A REGISTRATION NO. 279-2261

ACTIVE INGREDIENTS:

BY WT.

Q, Q-Dimethyl S-4-oxo-1,2,3-benzotriazin-	
3 (4 H) ylmethyl phosphorodithioate	- 3.00%
Sulphur	- - - - - 42.00%

INERT INGREDIENTS - - - - - 55.00%

† U.S. Patent No. 2,758,115

Canadian Patent No. 552,660

100.00%



DANGER - POISON



KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

USAGE CAUTION

Do not graze livestock in treated areas for 21 days after treatment.

Do not treat food crops grown in the green house.
Dispose of wastes by burying in non-crop land away from
water supplies.

Empty containers should be disposed of by burying with wastes or by burning. Keep Out of Smoke.

Highly toxic to fish. Do not contaminate any body of water nor apply to any area not specified on the label. Applications of materials containing Sulfur should not be made just prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Guthion is a registered trademark, Reg. U.S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G., Chemagro Corporation Licensee.

NEW - REVISED

ON TOMATOES to control Colorado Potato Beetle, Green Stinkbug, *Drosophila* and Leafminers use 15-25 pounds per acre. To control Aphids, European Corn borer, Flea Beetles, Hornworm, Leafhoppers and Thrips use 20-25 pounds per acre. To control Corn Earworm, Fruitworm, Pinworm, Tuberworm, and Yellow-striped Armyworm use 25 pounds per acre. No time limitations.

ON DRY ONIONS to control Thrips use 20-25 pounds per acre. Do not apply within 28 days of harvest. Do not apply more than 3 times per season.

ON ORNAMENTALS and NURSERY PLANTS to control Aphids, Lacebugs, Leafhoppers, Wites and Thrips use 20 pounds per acre. Dust all foliage surfaces including the underside of leaves for complete coverage. Do not allow children or pets on treated areas until material has been washed into soil and treated area is dry.

WARNING

Poisonous if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Wear protective clothing, natural rubber gloves, and goggles.

Do not breathe dust. Wear a mask or respirator effective for protection against GUTHION. Information as to suitable types of masks or respirators is available from the U.S. Department of Agriculture.

Do not contaminate feed or food. Keep out of reach of children. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Workers who must enter treated fields within 2 days of application should wear protective clothing. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

TREATMENT: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

ACCEPTED

MAR 13 1968

279-2261

Printed in U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



LABEL

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE - On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY - Niagara warrants that this product, as of the time it is sold by Niagara, conforms to the ingredient statement on the label and is reasonably fit for the purposes set forth on the label and for use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD "AS IS."** NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES - Buyer or user's exclusive remedy for damages in breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS - Because of certain critical conditions created or incurred by buyer or user, in which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK



FRENO, CALIF. - RICHMOND, CALIF.
RIVERSIDE, CALIF. - YAKIMA, WASHINGTON