

Page 3

PH Key No. 17-2001

Code 50202

Net Weight

Kolodust® 50 G-3 Dust

Contains 3% Guthion®

Insecticide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-226 EPA Est. 274

Active Ingredients	B. W.
O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3 (4H) yl)-methyl] phosphorodithioate	3.00%
Sulphur	42.00%
Inert ingredients	55.00%
	100.00%

Guthion® is a registered trademark. Reg. U.S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G. Chemagro Corporation Licensee.
Canadian Patent No. 552,661

KEEP OUT OF REACH OF CHILDREN




For Emergency Assistance Call 716-735-3765

See Other Panels for Additional Precautionary Information



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19102

PA 1-80



Symptoms of Poisoning—A sense of dizziness, headache, weakness, contracted pupils, blurred vision, vomiting and diarrhea.

First Aid

In case of poisoning, call a physician immediately. Have patient lie down and keep quiet.

If **swallowed**, drink plenty of glasses of water and induce vomiting by touching back of throat with finger. Do not attempt to induce vomiting if the patient is unconscious or if the patient is a child.

If on **skin**, remove contaminated clothing and wash skin immediately with soap and warm water.

If in **eyes**, wash with flowing water for at least 15 minutes.

Antidote

Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. Atropine is a secretory inhibitor and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema which may develop in serious cases of poisoning within 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Danger

Poisonous if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Wear protective clothing (goggles, gloves, and goggles).

Do not breathe dust. Wear a mask or respirator (effective for protection against Guthion) only as approved by the Mining Enforcement and Safety Administration, formerly the U.S. Bureau of Mines, and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash and decontaminate clothing with soap and hot water before reuse.

Keep all unprotected persons out of the operating area of equipment where there may be danger of drift. Workers who must enter treated fields within 2 days of application should wear protective clothing.

Environmental Hazards

Highly toxic to fish and wild life. Birds and other animals may be killed. Keep out of any body of water. Do not use in areas where cleaning of equipment or dusting of machinery is necessary. Conditions favor drift from the treated area. Do not apply to the area not specified on the label. Do not use on Melons, Squash, Cucumbers, Apples, and other fruits which tolerate sulphur. Applications of materials containing sulphur should not be made just prior to flowering periods. Do not apply at high temperatures nor within 10 days to 2 weeks before harvest.

Do not allow this material to come into contact with people or animals. Do not use in areas where children or animals may be present. Do not use in areas where label directions are not followed. Do not use in areas where other undesirable results may occur.

Page 2 of 3

DIRECTIONS FOR USE

Use in accordance with Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or material that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or by incineration. Place away from water supplies. Dispose of containers by incineration or landfill approved for pesticide containers. Do not use in a safe place. Consult Federal, State or local authorities for approved alternative disposal methods. See instructions for disposal.

Do not graze livestock in treated areas until the label indicates.

Do not treat food grown in a greenhouse.

Apply when pests or disease first appear and repeat as necessary. For control beyond the usage limitations of this product, use a suitable material. May be applied with airplane or ground application equipment. For Mildew control, the material must be applied before the disease appears. Observe days interval between applications and harvest indicated by number in following table.

Apples (15), Crabapples (15), Pears (15), Quinces (15): Aphids, Apple Maggot, Codling Moth, European Apple Sawfly, Eyespot, Black Moth, Fruit Tree Leafroller, Leafhoppers, Mites, Redbanded Leafhopper, Powdery Mildew. Use 10 to 20 pounds per acre. Do not apply more than 4 times per season.

Beans: Aphids, Bean Leafbeetle, Leafhoppers, Leafminers, Leafrollers, Mites, Sunk Bugs. Use 15 to 20 pounds per acre. Mexican Bean Beetle, Powdery Mildew, Spotted Cucumber Beetle, Striped Cucumber Beetle, Tarnished Plant Bug, Western Striped Cucumber Beetle. Use 20 pounds per acre. Do not apply to Snap Beans within 7 days of harvest. Do not apply to Dry Beans within 30 days of harvest. Do not apply more than 4 applications. Do not feed ensilage or treated forage. Sulphur may injure certain bean varieties, check tolerance prior to use.

Blackberries (14), Boysenberries (14), Loganberries (14), Raspberries (14): Aphids, Leafhoppers, Leafminers, Leafrollers, Sawflies, Root Weevil, Powdery Mildew. Use 2 to 3 pounds per acre.

Broccoli (15), Brussels Sprouts (7), Cabbage (21), Cauliflower (15): Aphids, Cabbage Looper, Diamond Borer, Imported Cabbageworm. Use 20 to 25 pounds per acre.

Cherries (15): Brown Bark Beetle, Cherry Bud Moth, Cherry Leafminer, Mites, Plum Curculionid, San Jose Scale, Silver Leaf Miner. Do not apply more than 4 times per season.

Cotton: Aphids, Boll Weevil, Bollworm, Brown Cotton Leafworm, Cotton Fleahopper, Cotton Leafworm, Lygus Bugs, Mites, Pink Bollworm, Rapid Plant Bug, Stink Bug, Tarnished Plant Bug, Thrips. Use 16 to 33 pounds per acre. Do not apply within 1 day of picking. If late season applications are made, do not graze livestock on treated areas or feed gin wastes.

Cowpeas (Southern Peas, Blackeyed Peas, Crowder Peas) (7): Leafminers, Stink Bugs—Use 15 to 33 pounds per acre. Corn Earworm, Cowpea Curculio—Use 33 pounds per acre. Do not apply more than 4 times per season. Do not use vines for feed or forage nor pasture treated areas. Sulphur may injure certain varieties; check tolerance prior to use.

Dry Onions (28): Thrips—Use 20 to 25 pounds per acre. Do not apply more than 3 times per season.

Grapes: Grape Berry Moth, Leafhoppers, Mites, Powdery Mildew, Thrips (California)—Use 20 to 30 pounds per acre. Do not apply within 10 days of harvest. Redbanded Leafroller—Use 25 to 30 pounds per acre. Do not apply within 28 days of harvest. Do not make more than 3 applications per season.

Nectarines (21), (Peaches (21): Aphids, Brown Rot, Mites, Oriental Fruit Moth, Peach Twig Borer (California), Powdery Mildew, Redbanded Leafroller, Variegated (Platyedra Flavedana) Leafroller—Use 35 to 80 pounds per acre. Do not apply more than 8 times per season.

Ornamentals, Nursery Plants: Aphids, Lacebugs, Leafhopper, Mites, Thrips—Use 20 pounds per acre. Dust all foliage surfaces including the underside of leaves for complete coverage. Do not allow children or pets on treated areas until material has been washed into soil and treated area is dry.

Plums (15), Prunes (15): Aphids, Brown Rot, E, spotted Bud Moth, Fruit Tree Leafroller, Mites, Peach Twig Borer (California), Redbanded Leafroller—Use 35 to 80 pounds per acre. Do not apply more than 8 times per season.

Potatoes (7): Colorado Potato Beetles, Leafminers—Use 15 to 25 pounds per acre. Aphids, European Cornborer, Flea Beetles, Leafhoppers, Spittlebugs, Tarnished Plant Bugs, Tuberworms—Use 20 to 25 pounds per acre.

Spinach (14): Aphids, Leafminers, Mites—Use 15 to 25 pounds per acre.

Tomatoes: Colorado Potato Beetle, Drosophila Greer, Stinkbug, Leafminers—Use 15 to 25 pounds per acre. Aphids, European Cornborer, Flea Beetles, Hornworms, Leafhoppers, Thrips—Use 20 to 25 pounds per acre. Corn Earworm, Fruitworm, Pinworm, Tuberworm, Yellow-striped Armyworm—Use 25 pounds per acre. No time limitations.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.