

Code: 31753



28 Gallons Net Contents

# PAR F 60 SOLUBLE OIL

### DIRECTIONS FOR USE

Apply when insects first appear and repeat as necessary. Do not use in undiluted form.

**LEMONS:** Red Scale, Black Scale, Red Spider -- Use 1.2-1.4 gallons per 100 gallons of water. Apply in late spring and late summer thru December.

**NAVEL AND VALENCIA ORANGES:** Red Scale, Yellow Scale, Citricola Scale, Purple Scale, Black Scale, Red Spider -- Use 1.2-1.4 gallons per 100 gallons of water. Apply in late summer thru October.

**PEACH, PLUM: Summer Use** Spider Mites -- Use 1-1 1/2 gallons per 100 gallons of water. Certain varieties of plums may be injured by oil sprays, check tolerance prior to treatment.

**APPLE, PEAR: Summer Use** Spider Mites -- Use 1-1 1/2 gallons per 100 gallons of water.

**PRUNES: Summer Use** Spider Mites -- Use 1-1 1/2 gallons per 100 gallons of water. Certain varieties of prunes may be injured by oil spray, check tolerance prior to treatment.

**OLIVES:** Olive Parlatoria Scale, Oleander Scale, Black Scale -- Use 1 1/2-2 gallons per 100 gallons of water. May be applied during the post-bloom through post-harvest periods.

**WALNUTS: Summer Use** Spider Mites, Soft Scales -- Use 1 gallon per 100 gallons of water. Apply 200 to 400 gallons of water per acre. Do not apply more than 4 gallons of Par F 60 Soluble Oil per acre. **Delayed Dormant** -- Scale, Overwintering Mite Eggs -- Use 2 gallons per 100 gallons of water. Apply in sufficient water for thorough coverage. Apply a minimum of 200 gallons of water per acre. This treatment should only be used on trees that have not suffered from lack of moisture at any time during the growing season. Further, oil should not be applied to orchards where soil moisture is low.

**ALMONDS, CHERRIES: Summer Use** Red Spider Mite -- Use 2-4 quarts per 100 gallons of water. To control summer mites of Brown Apricot Scale, San Jose Scale, 1 1/2-4 1/2 quarts per 100 gallons of water. **Dormant or delayed Dormant** -- Apricot Scale, Black Scale, San Jose Scale, Parlatoria Scale, Mite Eggs, Aphid Eggs -- Use 2 gallons per 100 gallons of water. **Delayed Dormant** -- Mite Eggs -- Use 2 1/2 gallons per 100 gallons of water.

ACCEPTED  
JUL 2 1961  
279-2292

### INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-2292 AA

ACTIVE INGREDIENT:	BY WT.
Petroleum Oil	99.00%
INERT INGREDIENTS:	1.00%
	100.00%

Classification -- Unclassified

Unsulforated Residue -- 96% Minimum

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Avoid inhalation of spray mist. Harmful if swallowed. Do not induce vomiting. Call a Physician immediately.

DO NOT USE THIS MATERIAL IF IT DOES NOT EMULSIFY

### MIXING DIRECTIONS

Before using this product, be sure the spray tank is clean and contains just enough water to cover the agitator shaft.

If an injector is used, first add 25 to 50 gallons of water to tank with engine running to provide good pressure. Then put injector suction tube into the measured amount of spray oil, opening injector valve to allow the oil to be sucked into the tank. Then fill the tank with water.

Use only in equipment with mechanical or paddle agitation. Be sure spray tank is clean. With enough water in tank to cover agitator shaft add desired amount of this material to the tank with agitator in motion. Pump this mixture through the overflow and back into tank for about two minutes. If this mixture turns white, it indicates good emulsification. Fill the tank with water, adding other desired materials as tank is filling. Keep agitator running at all times.

### USAGE CAUTIONS

Use efficient equipment of the proper type; avoid spraying during or immediately prior to hot or freezing weather, hot dry winds, rain or other unsuitable conditions; avoid overspraying or double spraying. Plants should be sprayed only when in vigorous condition and when their moisture condition is suitable.

On Deciduous trees do not mix with sulfur or apply within 10 days to 2 weeks prior to or after a sulfur treatment.

On Citrus do not mix with sulfur or apply within 60 days prior to or after a sulfur treatment.

Do not mix with pesticides incompatible with oil.

**DISPOSE OF WASTES** by burying in non-crop land away from water supplies. Empty 1 and 5 gallon cans should be smashed and buried with wastes. Empty drums should be returned to your pesticide supplier. Do not re-use.

Do not allow this material to drift onto neighboring crops or non crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Dealer Should Sell in Original Packages Only.

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:

**WARRANTY:** Niagara warrants that this product, as of the time of sale by Niagara, conforms to the ingredient statement on the label and is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

**DAMAGES:** Buyer's or user's exclusive remedy for claims for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 90 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

**EXCUSE FROM NON CONTROLLABLE CONDITIONS:** Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

NIAGARA CHEMICAL DIVISION  
FMC CORPORATION  
MIDDLEBURY, N.Y.

**fmc** CORPORATION  
FRESNO, CALIF. • RICHMOND, CALIF. • YAKIMA, WASHINGTON