

CODE  
20070

(Contra Parasit)  
SULFUR 60 BORON PEANUT DUST  
WITH BORAX  
Fungicide - Nutritional

NET WEIGHT  
50 LBS.

U. S. D. A. Reg. No. 279-2316

ACTIVE INGREDIENTS:

279 2316

\*Sulfur ----- 60.0%

\*\*Copper, expressed as metallic ----- 4.0%

INERT INGREDIENTS:\*\*\* ----- 36.0%

100.0%

\*Sulfur Fineness: 93% through 325 mesh

\*\*In copper oxide form

\*\*\*Boron expressed as metallic ----- 0.34%

**CAUTION**

KEEP OUT OF REACH OF CHILDREN

Harmful If Swallowed.



CODE  
20070

FLUOROSULFONIC ACID  
SULFUR & SULFUR BORON FERTILIZER  
(10% SULFUR)  
10-10-10-10-10-10-10

REG. NO. 10070  
U.S.

#### DIRECTIONS FOR USE

CROP: PEANUTS

PEST: Leafspot, aid in correcting the deficiency of boron.

Apply undiluted as a dust at the rate of 15 pounds per acre when spots first appear on old leaves at the base of the plants. Only one application should be made per season. Dusting should be done with a machine capable of depositing a thin uniform coating of dust on all above-ground plant surfaces. Dusting should only be done when the foliage is dry. Severe burning may result if application is made when the leaves are wet. Use of more than the recommended dosage may result in reduced yields.

Applications of dusting or spraying materials containing sulfur should not be made just prior to or during periods of excessively high temperatures.

CODE

COPIES 4 SULTUR 60 BORON PEANUT DUST

(WITH MAX)

## References

NET WELFARE

10 LBS.

## USAGE CAUTIONS

This product is toxic to fish. Keep out of any body of water.

Apply this product only as specified on this label.

Dispose of wastes by burying in non-crop lands away from water

supplies. Dispose of bags by burying with wastes or by burning.

Keep out of angle.

NOT ALL OF THIS MATERIAL TO BE DISCLOSED

1. The first group of authors (see Table 1) has shown that the use of a single, non-specific, and non-validated questionnaire to assess the prevalence of mental health problems in the general population is not sufficient to provide a reliable estimate of the prevalence of mental health problems. The use of a single questionnaire may also be insufficient to provide a reliable estimate of the prevalence of mental health problems in the general population.

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED  
DATE 07-16-2010 BY 60322 UCBAW

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY

FEBRUARY 1971  
 Volume 10, Number 2

Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D).

FIGURE 14-4

Environ Monit Assess (2008) 142:111–120  
DOI 10.1007/s10661-008-9400-2

[illegible][illegible]