

Code: 31055



50 Pounds

SULPHUR 25 SEVIN 10 DUST

DIRECTIONS FOR USE

Niagara Sulphur 25 Sevin 10 Dust may be applied with ground or airplane dusting equipment to give thorough coverage of plants with uniform deposit of dust. Apply when insects appear and repeat every 7-10 days if necessary. Apply thoroughly to upper and lower leaf surfaces. Sulphur must be applied before mildew appears for adequate control.

CROP	PEST	RATE AND REMARKS
Beans	Lygus Bugs Stink Bugs Leafhoppers Corn Earworm Western Bean Cutworm	Use 20 lbs. per acre. Apply when insects appear and repeat every 7 to 10 days if necessary. Sulphur may injure some varieties. Check tolerance before using.
Potatoes	Lygus Bugs Leafhoppers Flea Beetles Colorado Potato Beetle	Use 20 lbs. per acre. Apply when insects appear and repeat as necessary.
Grapes	Grape Leafhoppers Grape Leafhoppers Powdery Mildew	Use 20 lbs. per acre. For Leafhoppers apply when insects appear and repeat as necessary. For Leafhoppers, apply just before larvae emerge from first rolls. For second and third broods use Sevin alone or with smaller amounts of sulphur to prevent burn to the foliage.
Cotton	Bollworms Leaf Worms Flea Beetles Pink Bollworms Leaf Perforators	Use 20-25 lbs. per acre. Apply when insects appear and repeat as necessary.
Peaches	Peach Twig Borer Oriental Fruit Moth Catfacing Insects Powdery Mildew	Use 50-60 lbs. per acre. Do not apply within 1 day of harvest.

ACTIVE INGREDIENTS:	BY WT.
Carbaryl (1-naphthyl N-methylcarbamate)	10.00%
Sulphur	25.00%
INERT INGREDIENTS	65.00%
	100.00%

*Trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate (carbaryl)

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed.
 Avoid Prolonged or Repeated Breathing of dust.
 Do not take internally.
 Skin contact may be harmful.
 Avoid Unnecessary Contact. Wash hands and face before eating. Take shower or bath after work.
 Wear regular long sleeved work clothing. Change to clean clothing daily.
 Avoid contamination of Food, Feed, Feeding Troughs, and Watering Receptacles.
NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Manufactured in U.S.A. by

NIAGARA CHEMICAL COMPANY

FMC CORPORATION

ANAHEIM, CALIF. ● RICHMOND, CALIF.

FRESNO, CALIF. ● YAKIMA, WASH.

Home Office: Middleport, N. Y.



Applications of material should be made just prior to or during high temperatures.

Do not use on or allow on cucumbers, apricots or orange sulphur.

For protection of honeybees during periods when honeybees are necessary to use keepers well in advance until one week after.

CAUTION: Use of this product should be in accordance with label directions to avoid excessive residues or injury to crops.

Dealers Should Sell



50 Pounds Net Weight

SULPHUR 25 SEVIN*10 DUST

with ground coverage of insects apply thoroughly must be ap-

Apply when repeat every necessary. Sulphur varieties. before using.

Apply when repeat as nec

re. For Leaf- insects ap- necessary. apply just be- and third of sulphur to

acre. Apply and repeat

acre. Do not of harvest.

ACTIVE INGREDIENTS:	BY WT.
Carbaryl (1-naphthyl N-methylcarbamate) -	10.00%
Sulphur - - - - -	25.00%
INERT INGREDIENTS - - - - -	65.00%
	100.00%

*Trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate (carbaryl)

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed.
 Avoid Prolonged or Repeated Breathing of dust.
 Do not take internally.
 Skin contact may be harmful.
 Avoid Unnecessary Contact. Wash hands and face before eating. Take shower or bath after work.
 Wear regular long sleeved work clothing. Change to clean clothing daily.
 Avoid contamination of Food, Feed, Feeding Troughs, and Watering Receptacles.
NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Manufactured in U.S.A. by

NIAGARA CHEMICAL DIVISION

SMC CORPORATION

ANAHEIM, CALIF. ● RICHMOND, CALIF.

FRESNO, CALIF. ● YAKIMA, WASH.

Home Office: Middleport, N. Y.



USAGE CAUTION

Applications of materials containing Sulphur should not be made just prior to or during periods of excessively high temperatures.

Do not use on or allow to drift onto melons, squash, cucumbers, apricots or other plants which will not tolerate sulphur.

For protection of honey bees avoid use, if possible during periods when honey bees are visiting the crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

CAUTION: Use of this material in a manner or at a time other than in accordance with the directions may produce plant injury, excessive residues or other undesirable results.

Dealers Should Sell in Original Packages Only.

SULPHUR 25

25 Pounds Net Weight
DUST

With good coverage of insects. Apply before most insects.

e. Apply when repeat every necessary. Sub-varieties. pre using.

e. Apply when repeat as nec

For leaf insects. Apply to

Apply to

40% SULPHUR	25.00
10% SULPHUR	10.00
5% SULPHUR	5.00
1% SULPHUR	1.00
NET WEIGHT	25.00
GROSS WEIGHT	27.00

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed.
 Avoid Prolonged or Repeated Breathing of dust.
 Do not take internally.
 Skin contact may be harmful.
 Avoid Unnecessary Contact. Wash hands and face before eating. Take shower or bath after work.
 Wear regular long sleeved work clothing. Change to clean clothing daily.
 Avoid contamination of Food, Feed, Fruit and Watering Receptacles.

NOTE FOR PHYSICIAN: Carbaryl is a carbamate, inhibits cholinesterase activity. At 100 mg/kg.

Manufactured in U.S.A. by



ANAHEIM, CALIF. RICHMOND, CALIF.
 FRESNO, CALIF. FARMINGDALE, N.Y.
 Home Office: Middletown, N.Y.





PLEASE CAUTION:

For protection of honey bees avoid use, if possible during periods when honey bees are visiting the crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance. Do not spray for a week after application.

CAUTION: Use of this material in a manner or at a time other than in accordance with the directions may produce plant injury, excessive residues, or other undesirable results.

Buyers Should Seal in Original Packages Only.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE - On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY - Niagara warrants that this product, as of the time of sale by Niagara, to conform to the ingredient statement on the label and to be reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES - Buyer's or user's exclusive remedy for damages not covered by warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claims must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS - Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION

