

Code: 30423

[279-2329]

ETHION 4 SEVIN® 5 DUST

DIRECTIONS FOR USE

Apply when mites or insects appear and repeat as necessary. Full coverage of both upper and lower leaf surfaces are essential for good control. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in () following the crop.

BEANS (2)*: Mexican Bean Beetle — Use 10 pounds per acre. Bean Leaf Beetle, Twospotted Spider Mite, Thrips, Cucumber Beetles, Flea Beetles, Leafhoppers, Westernbean Cutworm, Armyworms, Corn Earworm, Stink Bugs, Tarnished Plant Bug — Use 20 pounds per acre. ***(4) California only:** Lygus Bugs, Bean Leaf Beetle, Lima Bean Pod Borer — Use 40 pounds per acre. Do not feed treated bean foliage to livestock.

GRAPES (15):** Grape Leafhopper, Grape Leafholder, Grape Leaf-skeletonizer, Twospotted Spider Mite, Willamette Mite — Use 20 to 30 pounds per acre. Make no more than 2 applications during the fruiting period. ***(30)** Make no more than 3 applications during the fruiting period. Do not apply to Salvador, Muscat or Fehr-Szago varieties of grapes as injury may occur. Ground application preferred.

PEACHES (30), NECTARINES (30): European Red Mite, Twospotted Spider Mite, Brown Mite — Use 40 to 60 pounds per acre. Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada — Use 60 pounds per acre. Codling Moth, Tussock Moth — Use 80 pounds per acre. For Oriental Fruit Moth and Tussock Moth control, apply at petal fall and every 10 to 14 days thereafter. Do not apply more than twice during the fruiting period.

PEPPERS (21): Serpentine Leafminer, Twospotted Spider Mite, Colorado Potato Beetle, Flea Beetles, Leafhoppers, Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug, Stink Bugs — Use 20 pounds per acre. Do not apply more than 3 times during the fruiting period.

PLUMS (45), PRUNES (45): Brown Mite, European Red Mite, Twospotted Spider Mite, Peach Twig Borer, Prune Leafhopper, Codling Moth, Orange Tortrix, Western Tussock Moth — Use 80 pounds per acre. Do not apply more than twice during the fruiting period. For Codling Moth control, apply at petal fall and every 10 to 14 days thereafter.

034W

INSECTICIDE-MITICIDE

E.P.A. REGISTRATION NO. 279-2329 AA

ACTIVE INGREDIENTS:	BY WT.
Ethion; 0,0,0',0'-Tetraethyl S,S'-methylene bisphosphorodithioate	4.00%
Carbaryl; 1-naphthyl N-methylcarbamate	5.00%
INERT INGREDIENTS:	91.00%
	<u>100.00%</u>

U.S. Patent No. 2,873,228.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. May be absorbed through skin. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Wash all contaminated clothing with soap and water before re-use. Avoid storage near or contamination of feed and foodstuffs.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE FOR PHYSICIANS: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

For Emergency Assistance Call
716-735-3765

STRAWBERRIES (2): Twospotted Spider Mite, Strawberry Leafroller, Strawberry Weevil.

TOMATOES (2): Tropical Spider Mite, Two miner, Colorado Potato Beetle, Flea Beetle, worm, Tomato Fruitworm, Tomato Hornworm, Stink Bugs — Use 20 pounds per acre.


ENVIRONMENTAL AND DISPOS

This product is toxic to fish and wild streams and ponds. Do not contaminate equipment or disposal of wastes. Do not conditions favor drift from the areas treated run-off is likely to occur. Apply this product the label.

This product is highly toxic to bees exposed or residues on crops. Protective information your Cooperative Agricultural Extension Service

DISPOSE OF WASTES by burying in non-water supplies. Empty bags should be buried. Keep out of smoke.

®Trademark of Union Carbide Corporation carbaryl.



EPA EST

129]

ETHION 4 SEVIN® 5 DUST

Net Weight

50 Pounds

USE FOR USE

Repeat and repeat as necessary. Full leaf surfaces are essential for good ground or air. Observe days interval harvest indicated by number in ()

— Use 10 pounds per acre. Bean Mite, Thrips, Cucumber Beetles, Flea and Cutworm, Armyworms, Corn Plant Bug — Use 20 pounds per acre. Do not feed treated bean

er, Grape Leafholder, Grape Leaf-Mite Willamette Mite -- Use 20 to more than 2 applications during the more than 3 applications during the Salvador, Muscat or Fehr-Szago occur. Ground application pre-

European Red Mite, Twospotted Mite to 60 pounds per acre. Oriental Periodical Cicada — Use 60 pounds Moth — Use 80 pounds per acre. Black Moth control, apply at petal fall or later. Do not apply more than twice

er, Twospotted Spider Mite, Colorado Leafhoppers, Fall Armyworm, Tomato, Tarnished Plant Bug, Stink Bugs apply more than 3 times during

Mite, European Red Mite, Two-spotted Mite, Prune Leafhopper, Codling Moth — Use 80 pounds per acre during the fruiting period. For control fall and every 10 to 14 days

INSECTICIDE-MITICIDE

E.P.A. REGISTRATION NO. 279-2329 AA

ACTIVE INGREDIENTS:	BY WT.
Ethion; 0,0,0',0'-Tetraethyl S,S'-methylene bisphosphorodithioate	4.00%
Carbaryl; 1-naphthyl N-methylcarbamate	5.00%
INERT INGREDIENTS:	91.00%
	<u>100.00%</u>

U.S. Patent No. 2,873,228.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. May be absorbed through skin. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Wash all contaminated clothing with soap and water before re-use. Avoid storage near or contamination of feed and foodstuffs.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE FOR PHYSICIANS: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

For Emergency Assistance Call
716-735-3765

STRAWBERRIES (2): Twospotted Spider Mite, Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil — Use 30 pounds per acre.

TOMATOES (2): Tropical Spider Mite, Twospotted Spider Mite, Leaf-miner, Colorado Potato Beetle, Flea Beetles, Leafhoppers, Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug, Stink Bugs — Use 20 pounds per acre.


ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

®Trademark of Union Carbide Corporation for the active ingredient carbaryl.



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.

274-2329



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals