

Code: 31526

50 Pounds Net Weight

SULPHUR 50 KELTHANE*4 SEVIN**10 DUST

DIRECTIONS FOR USE

Apply when insects or disease first appear and repeat as necessary. For insect or disease control beyond the time limitations of this product use a suitable material. May be applied with ground or airplane application equipment.

ON BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES and RASPBERRIES to control Leafrollers including Omnivorous Leafroller, Sawfly, Climbing Cutworms, Snowy Tree Cricket, Orange Tortrix, Leafhoppers, Aphids, and Powdery Mildew use 20 pounds per acre. Do not apply within 7 days of harvest.

ON GRAPES to control Powdery Mildew, Grape Leaf Folder, Grape Leafhopper, Grape Leaf Skeletonizer and Mites use 30 pounds per acre. Apply just before first brood leaf folder larvae emerge from rolls and as needed for Leafhoppers. Do not apply within 7 days of harvest. Use ground application equipment only.

ON STRAWBERRIES to control Powdery Mildew, Meadow Spittlebug, Strawberry Leafroller and Strawberry Weevil use 20 pounds per acre. Do not apply within 2 days of harvest.

ON BEANS to control Mexican Bean Beetle use 10-20 pounds per acre. Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Leafhoppers, Corn Earworm, Stink Bugs and Tarnished Plant Bug use 20 pounds per acre. California: Lygus Bug, Stink Bug and Lima Bean Pod Borer use 20 pounds per acre. Do not apply within 7 days of harvest. Do not feed treated vines to meat or dairy animals. Sulphur may cause injury to certain varieties. Check sulphur Tolerance prior to use.

ON POTATO, TOMATO, and PEPPERS to control Colorado Potato Beetle, Flea Beetles, Leafhoppers, use 10-20 pounds per acre. To control Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished plant bugs, Russet Mite and Stink Bugs use 20 pounds per acre. No time limitations on Potatoes. On Tomato and Peppers do not apply within 2 days of harvest.

ON SUGAR BEETS to control Armyworm, Leafhoppers, Grasshoppers and Weevils use 20 pounds per acre. Do not apply within 21 days of harvest. Do not feed crop residues (tops) to dairy or meat animals.

ON COTTON Early Season: Thrip, Flea Beetles, Cotton Flea hopper, Cotton Leafworm and Stripped Blister Beetle use 10-25 pounds per acre. Mid and Late Season: Boll Weevil, Bollworm, Fall Armyworm, Cotton Leaf Perforator, Leafrollers, Leafhoppers, Tarnished Plant Bug, Lygus Bug and Aphids (suppression) use 20-25 pounds per acre. To control Pink Bollworm use 20-25 pounds per acre. To control Stink Bug and Salt Marsh Caterpillar use 20-25 pounds per acre. To control Grasshoppers use 10-25 pounds. Do not apply within 14 days of harvest. Do not graze treated fields by meat or dairy animals.

ON PEACHES and NECTARINES to control Oriental Fruit Moth, Peach Twig Borer, Periodical cicada, Codling Moth, Tussock Moth and Mites use 40-80 pounds per acre. On Peaches do not apply within 14 days, on Nectarines do not apply within 14 days of harvest.

ON CHERRIES, PLUMS and PRUNES to control Cherry Fruit Fly, Peach Twig Borer, Leafhopper, Codling Moth, Orange Tortrix, Western Tussock Moth, Brown Rot, Powdery Mildew and Mites use 40-50 pounds per acre. Do not apply within 7 days of harvest. Do not reapply within 30 days.

ON FILBERTS to control Filbert Aphid, Filbert Leafroller, Filbert Moth and Mites use 40-50 pounds per acre. Do not apply within 14 days of harvest.

0674 - REVISED

FUNGICIDE-INSECTICIDE-MITICIDE

U.S.D.A. REGISTRATION NO. 279-2340

| ACTIVE INGREDIENTS: | BY WT. |
|---|---------|
| Carbaryl (1-naphthyl N-methylcarbamate) - - | 10.00% |
| Sulphur - - - - - | 50.00% |
| 1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol - - - - - | 4.00% |
| INERT INGREDIENTS - - - - - | 36.00% |
| | 100.00% |

**Trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate (carbaryl)

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with eyes or skin. Wash thoroughly after handling. Wash all contaminated clothing with soap and water before reuse. Avoid storage near feed and foodstuffs. Keep away from domestic animals, feed and foodstuffs.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

USAGE CAUTION


For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers, well in advance to locate hives at a safe distance until 1 week after application.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

Wastes should be disposed of by burying in non-crop land away from water supplies. Bags should be destroyed by burying with wastes or by burning. (Keep Out of Smoke)

Applications of materials containing sulphur should not be made just prior to or during periods of excessively high temperature or within 10 days to 2 weeks of an oil spray.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on the label.




ON ALMONDS to control Codling Moth, Filbert Worm, Fruit Tree Leafroller, Brown Rot and Mites use 40-80 pounds per acre. Do not apply within 14 days of harvest. Do not feed treated hulls to livestock. If hulls are to be fed, do not apply after nuts begin to form. **ON APPLES and PEARS** to control Powdery Mildew, Codling Moth, Green Apple Aphid, Apple Maggot, Red-banded Leaf roller, Pear Psylla, Periodical cicada and Mites use 40-80 pounds per acre. Do not apply within 7 days of harvest. Application within 30 days after full bloom may also provide apple thinning. To avoid this, delay use until at least 30 days after bloom.

*Trademark Rohm and Haas Co.

Made in U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE - On purchase or use of this product buyer and user agree to the following conditions.

WARRANTY - Niagara warrants that this product, as of the time of sale by Niagara, is conforming to the information statement on the label and is reasonably fit for the purposes set forth in the label directions. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED.

DAMAGES - Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 90 days of loss or injury by buyer or user stating which buyer or user wishes any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS - Release of non-controllable conditions created or incurred by buyer or user or user which Niagara has no control. Buyer and user, as a condition of purchase or use, assume responsibility for all release, Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT NEW YORK

