

Code: 31757

Net Weight  
50 Pounds

# BOTRAN<sup>(R)</sup> 6 DUST

## DIRECTIONS FOR USE

Use on the following crops at the recommended time of application and rate. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in ( ) following the crop.

**APRICOTS (1):** Fruit decay (Monilinia), Brown Rot (Rhizopus) — Use 50 pounds per acre. Apply 10 days and 1 day prior to harvest.

**BLACKBERRY (1), BOYSENBERRY (1), RASPBERRY (1) (Pacific Northwest Only):** Fruit Decay, (Botrytis Rot) — Use 50 pounds per acre. Use a maximum of 4 applications per season. Begin application just prior to bloom and continue at 7 day intervals.

**CHERRIES sweet (1):** Blossom Blight — Use 50 pounds per acre. Apply at Popcorn, Bloom, Full Bloom and Petal Fall. Fruit Decay (Monilinia), Brown Rot (Rhizopus) — Use 50 pounds per acre. Apply 10 days and 1 day prior to harvest.

**GRAPES (1):** Bunch Rot (Botrytis Rot), Storage Rot — Use 30 pounds per acre. Begin application while clusters are still loose and repeat application at 2 week intervals. Ground application preferred.

**HEAD LETTUCE (Field) (14):** Wilt (Botrytis Rot, Clerotinia) — Use 30 pounds per acre. Make first application at thinning. Repeat application 7 to 10 days later.

**LEAF LETTUCE (Greenhouse) (14):** Blight (Botrytis Rot) — Use 30 pounds per acre. Apply dust 7 days following transplanting. Repeat applications when plants are half mature. Do not apply to wilted plants or seedlings.

**ONIONS, GARLIC:** White Rot (Sclerotium) — Use 250 to 400 pounds per acre broadcast or 43,560 linear feet of furrow. Direct Seeding — Mix broadcast - incorporate) in top 1½ inches of soil 1 to 2 weeks before onions are seeded. Planting sets or buds — Apply as broadcast — Incorporation or in furrow with standard dusting equipment prior to planting sets or buds. Note: Plow to 8 inches and cross disc treated areas prior to seeding sensitive follow-up crops to prevent injury. Do not plant spinach as follow-up crop in treated soil.

**PEACHES (1), NECTARINES (1):** Brown Rot or Blossom Blight, (Monilinia) — Use 60 pounds per acre. Apply at pink bud and full bloom. Fruit decay, (Monilinia), Brown Rot (Rhizopus) — Use 60 pounds per acre. Apply 18 days, 10 days and 1 day prior to harvest.

**TOMATOES (Greenhouse) (0):** Stem Canker (Botrytis) — Use 30 pounds per acre. Dust stem of plant from ground level up to a height of 18 to 24 inches. Repeat weekly or as necessary.

**SNAP BEANS (2):** White Mold (Sclerotinia Sclerotiorum) — Use 50 pounds per acre on pole varieties. On bush varieties use 37 pounds per acre. Begin applications when disease is anticipated and continue at 7 day intervals during periods favorable to development of disease. Do not feed treated forage to livestock.

## FUNGICIDE

EPA REGISTRATION NO. 279-2386 AA

ACTIVE INGREDIENT:	BY WT.
*2,6-dichloro-4-nitroaniline	6.00%
INERT INGREDIENTS:	94.00%
	100.00%

\*Botran

## WARNING

### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin and eyes. In case of contact, remove contaminated clothing and wash skin with soap and water, for eyes flush with water and call a physician.

Do not use for green fruit.

Do not use on germinating seeds or annual seedlings except as suggested.

Do not feed treated forage to livestock.

**ROSE, HYDRANGEA:** Blight (Botrytis) — Use 30 pounds per acre. Field. Apply prior to shipment or storage. Stored: Thoroughly dust storage area and all canes and dead foliage; repeat application during long storage periods.

**CHRYSANTHEMUM, GERANIUM, ROSE:** Blight (Botrytis) — Use 30 pounds per acre. Stock cuttings or Green house plants — Begin when disease is anticipated or first appears. Dust foliage and flowers at 7 to 14 day intervals. During prolonged damp cool period apply every 5 to 7 days.

**GLADIOLUS:** Dry Rot, Stromatinia (Soil) — Use 250 pounds per acre as a broadcast treatment. Dust into furrow at planting time, so treated soils surrounds corms. Blight (Botrytis) — Use 30 pounds per acre. Dust thoroughly. Force fungicide between spikes. Apply to bunched cut flower spikes after harvest and before shipment or storage.

## ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on this label.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

© TUCO Products Company — Division of The Upjohn Company.  
U.S. Patent No. 3,119,736.

ACCEPTED

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER  
ED UNDER

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.

271 - 2380

LABEL

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by  
FMC CORPORATION  
AGRICULTURAL CHEMICAL DIVISION  
P.O. Box 1000, Lima, Peru

**FMC**  
Chemicals