

Code: 30068

50 Pounds

KOLODUST® 25 DIAZINON® 3 DUST

DIRECTIONS FOR USE

Begin applications when insects or disease first appear and repeat as necessary to maintain control. For disease control this material must be applied prior to disease development.

APPLY AS A DUST AT THE RATE OF 30 TO 40 POUNDS PER ACRE.

ON BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS and KALE for the control of Aphids, Diamond-Back Moth, Imported Cabbageworm and Powdery Mildew.

Do not apply on Broccoli or Cauliflower within 5 days; or on Cabbage within 7 days; or on Collards and Kale within 10 days of harvest.

ON TOMATOES for the control of Dipterous Leaf Miner, Vinegar Flies 'Drosophila', Aphids and Russet Mite. Do not apply within 3 days of harvest.

ON LIMA, POLE, and SNAP BEANS for the control of Aphids, Leaf Miners, Cucumber Beetles, Leafhopper 'Lima's', Powdery Mildew and Rust. Do not apply within 7 days of harvest. Sulphur may injure some varieties of beans, check sulphur tolerance prior to application.

ON STRAWBERRIES for the control of Aphids and Powdery Mildew. Do not apply within 5 days of harvest.

ON ONIONS to control Leafminers and Onion Thrips. Do not apply within 10 days of harvest.

APPLY AS A DUST AT THE RATE OF 33 POUNDS PER ACRE:

ON GRAPES to control Grape Leafhoppers, Grape Leaf Folder and Powdery Mildew. Do not apply within 10 days of harvest.


FUNGICIDE - INSECTICIDE

ACTIVE INGREDIENTS:	BY WT.
Sulphur - - - - -	21.00%
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate - - - -	3.00%
INERT INGREDIENTS - - - - -	76.00%
	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid storage near food and feed products. Wildlife in contact with treated areas may be harmed. Fish may be killed if their waters are contaminated with this product.



N.F.R.

USAGE C

Do not apply this material to melons, squash, cucumbers, as will not tolerate sulphur.

Applications of materials cont made prior to or during perio eratures.

Where disease is a problem of a suitable fungicide.

Diazinon is a registered tr Corporation for its patented No. 2,754,243.

USAGE CAUTION: Do not allow th neighboring crop or non-crop area time other than in accordance wil injury, excessive residues or ut occur.

Dealers Should Sell in Original

NOTICE: Niagara and Seller warrant, to the e product will conform to the ingredient statement on other warranty, express or implied covering t shall be held responsible in any manner for per loss of any kind resulting from the handling, st or not in accordance with directions, regardless Seller. Buyer assumes all risk and liability theref on these conditions. No person is authorized to this notice.