

KOLODUST[®] 25 DIAZINON[®] 3 DUST

DIRECTIONS FOR USE

Apply when insects or disease first appear and repeat as necessary to maintain control. For control beyond the use limitations of this product use a suitable material. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

FIGS (5) Vinegar Flea, Dried Fruit Beetles () Use 50-80 pounds per acre.

GRAPES (10) Grape Leafhoppers, Powdery Mildew () Use 33 pounds per acre.

STRAWBERRIES (5) Aphids, Powdery Mildew () Use 33-40 pounds per acre.

HOPS (14) Beet Armyworm, Corn Earworm, Climbing Cutworm, Aphids, Cucumber Beetle () Use 60 pounds per acre.

BROCCOLI (5), CABBAGE (7), CAULIFLOWER (5) Aphids, Diamond Borer, Mottled Leaf Miner, Cabbage Looper, Flea Beetles, Powdery Mildew () Use 33-50 pounds per acre.

COLLARDS (10), KALE (10), PARSLEY (12), SWISS CHARD (12), TURNIP GREENS (10) Aphids, Diamond Borer, Flea Beetles, Mottled Leaf Miner () Use 33-50 pounds per acre.

POLE (7), LIMA (7) and SNAP BEANS (7) Aphids, Leafhoppers, Cucumber Beetle () Use 33-45 pounds per acre. For control of certain varieties, check with your representative for use.

ONIONS (10) Trips, Flea Beetles, Ear Flies, Leaf Weevils () Use 40 pounds per acre.

CORN (10) Corn Earworm () Use 40-60 pounds per acre. Make first application when silks first appear. Make 2 or 3 repeat applications at 2 to 3 day intervals. Corn may be picked immediately following last application. Do not feed forage to livestock within 10 days following last application. Birds feeding in treated areas may die.

TOMATOES (3) Tobacco Leafworm, Vine Borers, Greenhorn, Aphids, Russian Mite () Use 33-50 pounds per acre.

ORNAMENTALS Aphids, Leafhoppers, White Flies, Flea Beetles, Powdery Mildew () Use 33-40 pounds per acre. For control of certain varieties, check with your representative for use.

LETTUCE (10) Aphids, Leafhoppers () Use 33-40 pounds per acre.

SPINACH (10) Aphids, Leafhoppers () Use 33-40 pounds per acre.

CELERY (10) Aphids, Flea Beetles () Use 33-40 pounds per acre.

BEETS (14) Aphids, Flea Beetles, Leafhoppers () Use 33 pounds per acre.

FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279-2422-AA

ACTIVE INGREDIENTS

0.0 Dimethyl 0/2 isopropyl 6-methyl 4-
pyridinyl phosphorothioate
Sulfur

INERT INGREDIENTS

BY WT

3.00%

21.00%

76.00%

100.00%

USAGE CAL

This product is toxic to fish and wildlife. Keep contaminated water by cleaning of equipment only as specified on the label.

This product is highly toxic to bees exposed to protective information must be obtained from Service.

Applications of materials containing sulfur, sh periods of excessively high temperatures.

Do not allow this material to be blown away.

Do not apply when weather conditions favor the when runoff is likely to occur.

DISPOSE OF WASTES: This product is not crop t bags should be burned with wastes or burned. K

For Emergency Ass
716-735-37

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with mucous membranes and clothing. Wash thoroughly after handling. Avoid storage near food and feed products.

This product contains Bentonite Sulfur. Bentonite Sulfur is made by the addition of fumed sulfur into Bentonite in the proportion of thirty parts sulfur to seventy parts Bentonite. A portion of the sulfur particles are water-insoluble (non-colloidal). The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, suspension in spray tank and safety to host plants.



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, N. Y. 14105