

Code: 31915

BOTRAN® 6 KOLODUST® 25 DUST

Net Weight
50 Pounds

DIRECTIONS FOR USE

For Powdery Mildew control apply when disease first appears and repeat as necessary. For other disease control see the directions below. May be applied with ground or airplane equipment. Observe days interval between last application and harvest indicated by number in following the crop.

GRAPES (1): Bunch Rot (Botrytis Rot), Storage Rot, Powdery Mildew — Use 30 pounds per acre. For Bunch Rot and Storage Rot control, begin application while clusters are still loose and repeat applications at two week intervals. Ground application equipment preferred.

PEACHES (1), NECTARINES (1): Brown Rot or Blossom Blight, (Monilinia), Powdery Mildew — Use 60 pounds per acre. Apply at pink bud and full bloom. Fruit decay, (Monilinia, Brown Rot, Rhizopus Rot, Powdery Mildew) — Use 60 pounds per acre. Apply 18 days, 10 days and 1 day prior to harvest.

SWEET CHERRIES (1): Blossom Blight — Use 50 pounds per acre. Apply at Pop Corn Blossom, Full Bloom, Petal Fall. Fruit decay, (Monilinia, Brown Rot, Rhizopus Rot) — Use 50 pounds per acre. Apply 10 days and 1 day prior to harvest.

LEAF LETTUCE (Green House) (14): Bight (Botrytis Rot) — Use 30 pounds per acre. Apply dust 10 days following transplanting. Repeat applications when plants are half mature. Do not apply to wilted plants or seedlings.

HEAD LETTUCE (14): Wilt (Botrytis Rot, Sclerotinia) — Use 30 pounds per acre. Apply first application at thinning. Repeat application 7 days later.

BLACKBERRY (7), BOYSENBERRY (7), RASPBERRY (7) (Pacific Northwest Only): Fruit decay, Botrytis Rot, Powdery Mildew — Use 50 pounds per acre. Use a maximum of 4 applications. Begin applications prior to bloom and continue at 7 day intervals.

SNAP BEANS (2): White Mold, Sclerotinia, Sclerotium — Powdery Mildew — Use 30 pounds per acre for pole varieties. The bush varieties use 37 pounds per acre. Begin applications when disease is anticipated and continue at 7 day intervals during periods favorable to development of disease. Do not feed treated forage to livestock. For any more certain varieties, look reference to sulfur prior to application.

FUNGICIDE

EPA REGISTRATION NO. 279-2498 AA

ACTIVE INGREDIENTS:	BY WT.
2,6-Dichloro 4-nitroaniline	6.00%
Sulphur	21.00%
INERT INGREDIENT:	73.00%
	100.00%

WARNING

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. In case of contact, remove contaminated clothing and wash skin with soap and water. For eyes, flush with water and call a physician.

USAGE CAUTIONS

Do not use for dried fruit.

Do not use on germinating seeds or annual seedlings except as suggested.

Do not feed treated forage to livestock.

This product is toxic to fish. Keep out of lakes, ponds, and streams. Do not contaminate water by washing of equipment or disposal of wastes. Apply this product only as specified on the label.

Applications of materials containing sulfur should not be made just prior to or during periods of excessively high temperatures, or within 10 days to 2 weeks of an oil spray.

Do not allow this material to drift onto sulfur sensitive crops.

Do not apply when weather conditions favor drift from the area treated. Do not apply when runoff is likely to occur.

DISPOSE OF WASTES: By burning — noncrop field away from water supplies. Empty bags should be buried with wastes or burned. Keep out of course.

Kolodust contains 11.0 percent Bentonite Sulfur. Bentonite Sulfur is made by the absorption of fused sulfur into Bentonite in the proportion of thirty parts sulfur and seventy parts Bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, suspensibility in spray tank and safety to host plants.

ACCEPTED

NOV 8 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR EUGENE POLSON REGISTERED
UNDER FLD 79-2498 SUBJECT
TO ATTACHED COMMENTS

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



7-7-59

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product, buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to buy or waive any statement contained herein.

For further information,
contact your local FMC
representative or write to:
FMC Chemicals, Inc.,
P.O. Box 1000, Houston, Texas 77001

FMC
Chemicals