

Code: 31918

Net Weight
50 Pounds

BOTRAN®* 6 SULPHUR 25 DUST

DIRECTIONS FOR USE

For Powdery Mildew control apply when disease first appears and repeat as necessary. For other disease control see the directions below. May be applied by ground or air. Observe days interval between application and harvest as indicated by number in () following the crop.

BLACKBERRIES (7), BOYSENBERRIES (7), RASPBERRIES (7) (Pacific Northwest Only): Fruit Decay (*Botrytis Rot*) and Powdery Mildew — Use 50 pounds per acre. Use a maximum of 4 applications. Begin application just prior to bloom and continue at 7 day intervals.

GRAPES (1): Bunch Rot, Storage Rot (*Botrytis Rot*) and Powdery Mildew—Use 30 pounds per acre. For Bunch Rot and Storage Rot control, begin application while clusters are still loose and repeat applications at two week interval. Ground application preferred.

HEAD LETTUCE (14): (Field) Wilt (*Botrytis Rot*, *Sclerotinia*) — Use 30 pounds per acre. Apply first application at thinning. Repeat application 7 days later.

LEAF LETTUCE (Greenhouse): Blight, (*Botrytis Rot*) — Use 30 pounds per acre. Apply dust 10 days following transplanting. Repeat applications when plants are half mature. Do not apply to wilted plants, seedlings or later than 14 days before harvest.

PEACHES, NECTARINES: Brown Rot, Blossom Blight, (*Monilinia*), and Powdery Mildew — Use 60 pounds per acre. Apply at pink bud and full bloom. Fruit decay (*Monilinia*), Brown Rot (*Rhizopus*) and Powdery Mildew — Use 60 pounds per acre. Apply 18 days, 10 days and 1 day prior to harvest.

SNAP BEANS (2): White Mold (*Sclerotinia Sclerotiorum*) and Powdery Mildew — Use 50 pounds per acre on pole varieties. On bush varieties use 37 pounds per acre. Begin applications when disease is anticipated and continue at 7 day intervals during periods favorable to development of disease. Do not feed treated forage to livestock. Sulphur may injure certain varieties; check tolerance to sulphur prior to application.

SWEET CHERRIES: Blossom Blight — Use 50 pounds per acre. Apply at Pop Corn, Bloom, Full Bloom and Petal Fall. Brown Rot, Fruit Decay (*Monilinia*), Rhizopus Rot (*Rhizopus*) — Use 50 pounds per acre. Apply 10 days and 1 day prior to harvest.

FUNGICIDE

EPA REGISTRATION NO. 279-2518 AA

ACTIVE INGREDIENTS:	BY WT.
*2,6-Dichloro-4-nitroaniline	6.00%
Sulphur	25.00%
INERT INGREDIENT:	69.00%
	100.00%

USAGE CAUTIONS

Do not use for dried fruit.

Do not use on germinating seeds or annual seedlings except as suggested.

Applications of materials containing sulphur should not be made just prior to or during periods of excessively high temperatures or within 10 days to 2 weeks of an oil spray.

Do not feed treated forage to livestock.

WARNING

KEEP OUT OF REACH OF CHILDREN


Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin and eyes. In case of contact, remove contaminated clothing and wash skin with soap and water, for eyes, flush with water and call a physician. Do not contaminate or store near feed or foodstuffs.

For Emergency Assistance Call
716-735-3765

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds, and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on this label. DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

©TUCO Products Company — Division of The Upjohn Company.



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



279-2518

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.



Manufactured In U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals