

Code: 31054



50 Pounds Net Weight

SULPHUR 20 ETHION 4 DUST

DIRECTIONS FOR USE

Apply when the disease, mites or insects appear and repeat as necessary. For disease control beyond the time limitations of this product use a suitable fungicide. Full coverage of both upper and lower leaf surfaces are essential to good control. May be applied with ground or airplane application equipment.

1. **ON COTTON, SEED ALFALFA and SEED CLOVER** to control Spider Mites, and Lygus Bugs use 25-40 pounds per acre. Do not apply on cotton after bolls begin to open. Cotton that has been treated with Ethion should not be grazed or fed to dairy animals or other livestock. Do not feed treated forage threshings or hullings to livestock. Workers who enter a treated cotton field within 24 hours should wear protective clothing.

2. **ON STRAWBERRIES** to control Two-Spotted Spider Mite, Powdery Mildew, and Strawberry Leafroller use at the rate of 30 pounds per acre. Thoroughly dust both leaf surfaces and down into the crown of the plants. Do not apply within 2 days of picking.

3. **ON PEACHES** to control European Red Mite, Two-Spotted Mite, Brown Almond Mite, Brown Rot and Powdery Mildew use 40-60 lbs. per acre. Do not apply more than 2 applications during the fruiting period. Do not apply within 30 days of harvest.

4. **ON GRAPES** to control Grape Leafhopper, Two-Spotted Mite, Willamette Mite and Powdery Mildew use at the rate of 20-30 lbs. per acre. Make no more than 2 applications within 15 days of harvest, or 3 applications within 30 days of harvest.

5. **ON BEANS** to control Bean Leaf Beetle, Two-Spotted Spider Mite, Powdery Mildew, Rust and Thrips use 20 lbs. per acre. Cover the lower leaf surfaces. Do not apply within 2 days of harvest. Do not feed treated foliage to livestock. To control Lygus Bugs, Bean Leaf Beetle, and Thrips, Two-Spotted Mite and Powdery Mildew use 30 to 40 lbs. per acre. Do not apply within 4 days of harvest. Do not feed treated foliage to livestock. Sulphur may injure some varieties. Check sulphur tolerance prior to use.

6. **ON TOMATOES** to control Tropical Mite, Two-Spotted Mite, Russet Mite and Leaf Miner use at the rate of 20 lbs. per acre. Do not apply within 2 days of harvest.

7. **ON ALMONDS** to control Brown Almond Mite, European Red Mite, Two-Spotted Mite and Brown Rot use 40-50 pounds per acre. Do not apply more than one application during the fruiting period. Do not apply within 30 days of harvest.

8. **ON APPLES** to control Brown Almond Mite, European Red Mite, Two-Spotted Mite and Powdery Mildew use 40-50 pounds per acre. Do not apply more than 3 applications during the fruiting period. Do not apply within 30 days of harvest. Sulphur may injure some varieties, check sulphur tolerance prior to use.

9. **ON PLUMS and PRUNES** to control Brown Almond Mite, European Red Mite, Two-Spotted Mite, Brown Rot and Prune Rust use 40 pounds per acre. Do not make more than 3 applications during the fruiting period. Do not apply within 21 days of harvest.

10. **ON PEPPERS** to control Serpentine Leaf Miner and Two-Spotted Mite use 30 lbs. per acre. Do not make more than 3 applications during the fruiting period. Do not apply within 21 days of harvest.

057W

FUNGICIDE - PESTICIDE


ACTIVE INGREDIENTS:	By Wt.
Ethion	4.00%
Sulphur	20.00%
INERT INGREDIENTS	76.00%
	100.00%

10, 0, 0, 0'-Tetraethyl S, S'-methylene bisphosphorodithioate.
U. S. Pat. No. 2,873,228.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of or storage near feed and foodstuffs.



UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
OR ECONOMIC PESTICIDE REGISTERED
UNDER NO. 2-2572

with Comments

Manufactured in U.S.A. by
NIAGARA CHEMICAL DIVISION
FMC CORPORATION



ANAHEIM, CALIF. ● RICHMOND, CALIF.
FRESNO, CALIF. ● YAKIMA, WASH.
Home Office: Middletown, N. Y.



USAGE CAUTION!

Applications of materials containing Sulphur should not be made just prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Do not apply this material to or allow it to drift onto melons, squash, cucumbers, apricots or other plants which will not tolerate sulphur.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

Dispose of wastes by burying in non-crop land away from water supplies. Empty containers should be disposed of by burying with wastes or by burning. Keep Out of Smoke.

USAGE CAUTION. Do not allow this material to drift onto neighboring crop or non-crop areas or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

Dealers Should Sell in Original Packages Only.

NOTICE: Niagara and Seller warrant, to the extent of the purchase price, that the product will conform to the ingredient statement set forth on the label but they make no other warranty. Niagara and Seller shall be held harmless for any loss of any kind or not in accordance with the product label. Seller, Buyer assumes all risk and liability therefrom and accepts and uses the product on these conditions. No person is authorized to vary or waive anything contained in this notice.

Currently being revised

Printed in U.S.A.

Code: 31054



50

SULPHUR 20 ETHION 4 DUST

DIRECTIONS FOR USE

Apply when the disease, mites or insects appear and repeat as necessary. For disease control beyond the time limitations of this product use a suitable fungicide. Full coverage of both upper and lower leaf surfaces are essential to good control. May be applied with ground or airplane application equipment.

ON COTTON, SEED ALFALFA and SEED CLOVER to control Spider Mites, and Lygus Bugs use 25-30 pounds per acre. Do not apply on cotton after bolls begin to open. Cotton that has been treated with Ethion should not be grazed or fed to dairy animals or other livestock. Do not feed treated forage to livestock. Workers who enter a treated cotton field within 24 hours should wear protective clothing.

ON STRAWBERRIES to control Two-Spotted Spider Mite, Powdery Mildew, and Strawberry Leafroller use at the rate of 30 pounds per acre. Thoroughly dust both leaf surfaces and down into the crown of the plants. Do not apply within 2 days of picking.

ON PEACHES to control European Red Mite, Two-Spotted Mite, Brown Almond Mite, Brown Rot and Powdery Mildew use 40-60 lbs. per acre. Do not apply more than 2 applications during the fruiting period. Do not apply within 30 days of harvest.

ON GRAPES to control Grape Leafhopper, Two-Spotted Mite, Willow Leaf Mite and Powdery Mildew use at the rate of 20-30 lbs. per acre. Make no more than 2 applications within 15 days of harvest, or 3 applications within 30 days of harvest.

ON BEANS to control Bean Leaf Beetle, Two-Spotted Spider Mite, Powdery Mildew, Rust and Thrips use 20 lbs. per acre. Cover the lower leaf surfaces. Do not apply within 2 days of harvest. Do not feed treated foliage to livestock. To control Lygus Bugs, Bean Leaf Beetle, Rust, Thrips, Two-Spotted Mite and Powdery Mildew use 30 to 40 lbs. per acre. Do not apply within 4 days of harvest. Do not feed treated foliage to livestock. Sulphur may injure some varieties. Check sulphur tolerance prior to use.

ON TOMATOES to control Tropical Mite, Two-Spotted Mite, Russet Mite and Leaf Miner use at the rate of 20 lbs. per acre. Do not apply within 2 days of harvest.

ON ALMONDS to control Brown Almond Mite, European Red Mite, Two-Spotted Mite and Brown Rot use 40-50 pounds per acre. Do not apply more than one application during the fruiting period. Do not apply within 30 days of harvest.

ON APPLES to control Brown Almond Mite, European Red Mite, Two-Spotted Mite and Powdery Mildew use 40-50 pounds per acre. Do not apply more than 3 applications during the fruiting period. Do not apply within 30 days of harvest. Sulphur may injure some varieties. Check sulphur tolerance prior to use.

ON PLUMS and PRUNES to control Brown Almond Mite, European Red Mite, Two-Spotted Mite, Brown Rot and Prune Rust use 40 pounds per acre. Do not use more than 3 applications during the fruiting period. Do not apply within 21 days of harvest.

ON PEPPERS to control Serpentine Leaf Miner and Two-Spotted Mite use 30 lbs. per acre. Do not use more than 3 applications during the fruiting period. Do not apply within 21 days of harvest.

FUNGICIDE - PESTICIDE

ACTIVE INGREDIENTS: Ethion 4, Sulphur 20
INERT INGREDIENTS: 76
NET WEIGHT 100 LBS.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Hazardous if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contact with food or storage near feed and foodstuffs.

REGISTERED

Manufactured in U.S.A. by
**NIAGARA CHEMICAL DIVISION
FMC CORPORATION**



ANAHEIM, CALIF. • RICHMOND, CALIF.
FRESNO, CALIF. • YAKIMA, WASH.
Home Office: Middleport, N. Y.



Applications of material just prior to or during harvest within 10 days to two weeks.

Do not apply this material on squash, cucumbers or melons.

Do not apply this material on plants that are stressed.

Dispose of wastes by burying in soil. Empty containers should be disposed of by burning.

UNSAFE: ETHION is highly toxic to fish and aquatic life. Do not use in or near streams, rivers, or other bodies of water.

Dealers Should Sell: This product is sold only to dealers who are registered with the manufacturer. Dealers should check the registration number on the label of this product.

0574



50 Pounds Net Weight

SULPHUR 20 ETHION 4 DUST

FUNGICIDE - PESTICIDE

ACTIVE INGREDIENTS
Ethion
Sulphur
INERT INGREDIENTS

By #1
4 004
2 004
76 004
100 004

Contains Tetraethyl Sulphur and Diethylphosphorothioate
U.S. Pat. No. 2,873,218

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of or storage near feed and foodstuffs.

ACCEPTED

With *[Signature]*

Manufactured in U.S.A. by
NIAGARA CHEMICAL DIVISION
FMC CORPORATION

ANAHEIM, CALIF. ● RICHMOND, CALIF
FRESNO, CALIF ● YAKIMA, WASH
Home Office: Middletown, N. Y.



USAGE CAUTIONS

Applications of materials containing Sulphur should not be made just prior to, or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Do not apply this material to or allow it to drift onto melons, squash, cucumbers, apricots or other plants which will not tolerate sulphur.

Keep out of reach of children. Keep out of reach of animals. Apply to plants only.

Dispose of wastes by burying in non-crop land away from water. Empty containers should be disposed of by burying with wastes or by burning. Keep Out of Smoke.

Warning: A toxic substance which may be harmful if inhaled. Avoid drift onto crops. Do not use on crops or areas of use in a manner which may cause the plant to be damaged with direct or indirect application. Excessive residues or other undesirable results may occur.

Dealers Should Sell in Original Packages Only

This product is sold in accordance with the terms of the license granted by the U.S. Department of Agriculture to the FMC Corporation. The ingredients are listed on the label. The label is to be read and followed carefully. The label is to be kept on the container at all times. The label is to be read and followed carefully. The label is to be kept on the container at all times.

Made in U.S.A.

Repeat and repeat as necessary. Applications of this product should be made to both upper and lower surfaces. May be applied with...
Control Spider Mites. Do not apply on cotton when treated with Ethion or other livestock. Do not apply to livestock. Workers should wear protective clothing.
Control Mite, Powdery Mildew, etc. 30 pounds per acre. Apply to the crown of the plant.
Control Spotted Mite, Brown Mite, etc. 40-60 lbs. per acre during the fruiting period.
Control Spotted Mite, Willamette Mite, etc. 30 lbs. per acre. Apply to the crown of harvest, or 3 weeks before harvest.
Control Spotted Spider Mite, etc. per acre. Cover the crown of harvest. Do not apply to Lygus Bugs, Bean Leaf Miner, etc. use 30 to 40 lbs. per acre. Do not feed to some varieties.
Control Spotted Mite, Russet Mite, etc. per acre. Do not apply to some varieties.
Control European Red Mite, Two-spotted Mite, etc. per acre. Do not apply to some varieties.
Control European Red Mite, Two-spotted Mite, etc. per acre. Do not apply to some varieties.
Control European Red Mite, etc. 40 pounds per acre. Do not apply to some varieties.
Control European Red Mite, etc. 40 pounds per acre. Do not apply to some varieties.