

Code: 31963

50 Pounds Net Weight

CAPTAN 10 KELTHANE* 4 DUST

DIRECTIONS FOR USE

Apply when pests or disease first appear and repeat as necessary. May be applied with airblast or ground application equipment.

APPLY TO GIVE THOROUGH COVERAGE TO UPPER AND LOWER LEAF SURFACES.

ON POTATOES to control Early Blight, Late Blight and Mites use 25-30 pounds per acre.

ON TOMATOES to control Early and Late Blight, Anthracnose, Septoria Leaf Spot and Mites. On young plants use 20 to 25 pounds per acre. On mature plants use 30 to 35 pounds per acre. Do not apply within 2 days of harvest.

ON STRAWBERRIES to control Botrytis, Grey Mold and Mites use 30 to 35 pounds per acre. Do not apply within 2 days of harvest.

ON APPLES to control Apple Scab, Brooks Fruit Spot, Fly-Specking, Bitter Rot, Black Rot, Botrytis, Black Pox, Clover, European Red and Two Spotted Mites use 40-50 pounds per acre. Do not apply within 7 days of harvest.

ON PEARS to control Pear Scab (except Far East Northwest), Clover, European Red and Two Spotted Mites use 40-50 pounds per acre. Do not apply within 7 days of harvest. Do not use on Udo or Pear Rust.

ON PEACHES to control Brown Rot, Jacket Rot, Two Spotted Mites and Spotted Mites use 50 pounds per acre. Do not apply within 14 days of harvest.

ON PLUMS AND PRUNES to control Brown Rot, Two Spotted Mites and Spotted Mites use 40-50 pounds per acre. Do not apply within 7 days of harvest. Do not apply within 30 days of the preceding application.

ON CHERRIES to control Cherry Leaf Spot, Brown Rot, Two Spotted Mites and Spotted Mites use 50-55 pounds per acre. Do not apply within 7 days of harvest. Do not repeat application within 30 days of the preceding application.

ON SWEET POTATOS, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELONS, PEPPERS, EGGSPLAINS, SWEET CORN, SWEET POTATOS, SWEET POTATOS, SWEET POTATOS to control Brown Rot, Two Spotted Mites and Spotted Mites use 20-25 pounds per acre. Do not apply within 2 days of harvest.

ON PEACHES AND NECTARINES to control Brown Rot, European Red, Two Spotted Mites and Spotted Mites use 50 pounds per acre. Do not apply within 14 days of harvest.

ON ALMONDS to control Brown Rot, Shot Hole, Green Leaf Blight, Clover, European Red and Two Spotted Mites use 40-50 pounds per acre. Do not apply within 14 days of harvest. Do not feed Almond Nuts to dairy animals or animals being fed to their children.

ON GRAPES to control Two Spotted Mite, European Mite, White Mite, Mite and Botrytis use 25-30 pounds per acre. Do not apply within 2 days of harvest.

To control European Red, starting July 1 to July 15, apply 4 dusts at 7 day intervals. Apply over the top and direct toward the center. The dusts are not to be applied with the wind. The dusts are not to be applied before rain. We ever rain, therefore, apply at least a little before rain. Do not apply following rain. Do not apply within 2 days of harvest. Use ground application equipment only.

FUNGICIDE - MITICIDE
U.S.P.A. REGISTRATION NO. 279 2567

ACTIVE INGREDIENTS:	BY WT.
7-Captan	10.00%
1,1-Dimethyl-2-propenyl-2,2,2-trichloroethyl	8.00%
INERT INGREDIENTS	86.00%
	100.00%

N-((trichloroethyl) (2,2,2-trichloroethyl)-1,2-dicarbonyl)

U.S. Patent Nos. 2,550,770 - 2,553,771 - 2,555,776

USAGE CAUTION

Do not use to fish. Keep out of the body of water. Do not apply when weather conditions favor drift from crop area treated.

Use a suitable fungicide where additional use are necessary for disease control beyond the usage instructions of this product.

Dispose of waste by burying in earth, away from water supplies. Bins and containers disposed of by burying with waste or by burning. Keep out of fire.

CAUTION KEEP OUT OF REACH OF CHILDREN


Avoid contact with dust. Avoid contact with skin. Avoid contact with eyes. Avoid contact with clothing. Avoid contact with food. Avoid contact with water. Avoid contact with animals. Avoid contact with children. Avoid contact with birds. Avoid contact with bees. Avoid contact with aquatic life. Avoid contact with fish. Avoid contact with shellfish. Avoid contact with mollusks. Avoid contact with crustaceans. Avoid contact with insects. Avoid contact with arachnids. Avoid contact with nematodes. Avoid contact with fungi. Avoid contact with bacteria. Avoid contact with viruses. Avoid contact with parasites. Avoid contact with weeds. Avoid contact with grasses. Avoid contact with shrubs. Avoid contact with trees. Avoid contact with crops. Avoid contact with livestock. Avoid contact with poultry. Avoid contact with swine. Avoid contact with horses. Avoid contact with dogs. Avoid contact with cats. Avoid contact with rabbits. Avoid contact with guinea pigs. Avoid contact with hamsters. Avoid contact with mice. Avoid contact with rats. Avoid contact with squirrels. Avoid contact with chipmunks. Avoid contact with chipmunks. Avoid contact with chipmunks.

*Trademark Rohm and Haas Company

11/10/1967

279-2567 U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers should See Original Package Only

For information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.

For more information on this product, contact your local FMC representative or write to FMC Corporation, Niagara Chemical Division, Middleport, New York.



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

