

Code: 31983

CAPTAN 5 SULPHUR 30 DUST 50 Pounds Net Weight

DIRECTIONS FOR USE

- May be applied with ground or airplane application equipment.
- ① ON PEACHES and NECTARINES: For control of Brown Rot, Powdery Mildew and Roscan Spot use 50-60 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.
 - ② ON PLUMS and PRUNES: to control Brown Rot, and Prune Rust use 50-60 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.
 - ③ ON STRAWBERRIES: For control of Grey Mold, and Powdery Mildew use 40-50 pounds per acre. Apply at first sign of disease. Repeat at weekly intervals as necessary to maintain control.
 - ④ ON TOMATOES to control Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot and Russet Mite - Young Plants 25 to 30 pounds per acre. Mature plants 40-50 pounds per acre. Begin applications when disease or mites first appear and repeat at 7-10 day intervals as necessary to maintain control.
 - ⑤ ON POTATOES to control Early Blight, Late Blight. Begin applications when plants are 2 to 4 inches high and repeat at 7-10 day intervals as long as disease threatens. Apply 50-55 pounds per acre, increasing to 40-50 pounds per acre on mature plants.
 - ⑥ ON CARNATIONS: Rose-Black Spot, Powdery Mildew and Rust. Carnations - Powdery Mildew, Rust and Leaf Spot - Chrysanthemums - Powdery Mildew and Rust - Apply at first sign of disease and repeat at 7-10 day intervals as necessary. Dust for complete coverage.

ACCEPTED

AUG 22 1967

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENT CONTROL ACT FOR ECONOMIC POISON, REGISTERED UNDER NO. 214-2296 SUBJECT TO ATTACHED COMMENTS.

FUNGICIDE

U. S. D. A. Reg. No. 279-

ACTIVE INGREDIENTS:	BY WT.
Captan - - - - -	5.00%
Sulfur - - - - -	50.00%
INERT INGREDIENTS - - - - -	45.00%
	100.00%
[Trichloromethyl] thio]-4-cyclohexene-1,2-dicarboximide U.S. Pat. Nos. 2,553,770 - 2,553, 771 2,553,776	

USAGE CAUTION

Applications of materials containing Sulphur should not be made just prior to or during periods of unusually high temperatures or within 10 days to two weeks of an oil spray.

Wash spray equipment and containers and dispose of wastes in non-crop land away from water supplies. Empty containers should be disposed of by burying with wastes or by burning. Keep Out of Smoke.

Fish may be killed if their waters are contaminated with this product. Do not apply when weather conditions favor drift from crop area treated.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid inhalation of dust. Avoid contact with skin.
Avoid contamination of feed or bedding.

Dealers Should Sell in Original Packages Only.

USAGE CAUTION: Do not allow this material to drift into neighboring crop or non-crop areas or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

TERMS OF SALE OR USE of this product shall be governed by the contract between the buyer and the seller.

WARRANTY: This product is warranted to be free from mechanical defects in the container and to contain the amount of active ingredient stated on the label. It is not warranted to be effective against all pests or diseases which it is claimed to control. It is not warranted to be effective under all conditions of use or in all localities.

INFORMATION: This product is a fungicide and should be used only on plants and crops for which it is labeled. It is not to be used on food crops unless specifically labeled for such use. It is not to be used on lawns, lawns, or other areas where it may be harmful to plants or animals. It is not to be used in or near water bodies.

PRECAUTIONS: This product is a dust and should be used with care to avoid inhalation of dust. It should be used in a well-ventilated area. It should not be used in or near water bodies. It should not be used in or near lawns, lawns, or other areas where it may be harmful to plants or animals.

Dust in U.S.A.