

ACCEPTED

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. V SUBJECT
TO ATTACHED COMMENTS.

Code: 31931



Gallons Net Contents

METHYL PARATHION 1.0 THIODAN® 2.0 C.O. E.C.

DIRECTIONS FOR USE

Apply when insects first appear and repeat as necessary to maintain control. Do not apply this product to plants in bloom. Use the maximum recommended rate of sufficient water to thoroughly cover plants. Broadcast applications require 10 gallons when applying this material by aircraft. Mix the recommended amount with sufficient water to provide a minimum of 1 gallon of spray per acre. Observe days interval between last application and harvest. Do not harvest any fruits or vegetables following the above.

ARTICHOKE (7): Apply 1 1/2 to 2 quarts per acre. Artichoke Plume Moth. Use 2 quarts per acre.

BEANS, Succulent and Dry (except Lima Beans) (15): Black Bean Aphid, Cucumber Beetle, Leafminer, Bean Leaf Skeletonizer, Green Stem Weevil, etc. Use 1 1/2 to 2 quarts per acre. Apply every 7 to 10 days. Do not exceed 3 applications per season. Do not feed treated thrashings to livestock or allow to graze in treated fields.

BROCCOLI (7), BRUSSELS SPROUTS (14), CAEBAGE (21), CAULIFLOWER (14): Aphid, Cabbage Looper, Imported Cabbageworm, Diamond-back Moth Larvae, Cross-striped Cabbageworm, Flea Beetle, Strik, Brass Hare, etc. Use 1 1/2 to 2 quarts per acre.

CELERY (Whole Plant) (15): Apply when insects first appear. Aphid, Cabbag Looper, Flea Beetle, etc. Use 1 1/2 to 2 quarts per acre. Apply every 7 to 10 days. Do not exceed 3 applications per season.

KALE (21): Apply when insects first appear. Flea Beetle, etc. Use 1 1/2 to 2 quarts per acre. Do not exceed 3 applications per season.

LETTUCE (21): Cabbage Looper, Aphid, Imported Green Flea Beetle, etc. Use 1 1/2 to 2 quarts per acre. Apply every 7 to 10 days. Do not exceed 3 applications per season.

PEAS, SUCCULENT (Northwest Only) (15): Apply when insects first appear. Flea Beetle, etc. Use 1 1/2 to 2 quarts per acre. Do not exceed 3 applications per season.

PEPPERS (15): Apply when insects first appear. Flea Beetle, etc. Use 1 1/2 to 2 quarts per acre. Do not exceed 3 applications per season.

SPINACH (21): Apply when insects first appear. Flea Beetle, etc. Use 1 1/2 to 2 quarts per acre. Do not exceed 3 applications per season.

TOMATOES (10): Apply when insects first appear. Flea Beetle, etc. Use 1 1/2 to 2 quarts per acre. Do not exceed 3 applications per season.

INSECTICIDE

U.S.D.A. REGISTRATION NO. 27-2609

ACTIVE INGREDIENTS	Per Cent
1.00 D.M.P. (1.0 parathion) methyl phosphorothioate	10.0
1.00 Endosulfan (1.0 thiodan) ethion	10.0
2.00 Inert ingredients	20.0
Petroleum Distillate	16.0
NET INGREDIENTS	56.0
	100.0

Contains 1.00 lb. of active ingredient per gallon. 2 p.p.m. of DTPP per gallon.



DANGER - POISON

KEEP OUT OF REACH OF CHILDREN



FIRST AID

If swallowed: Do not induce vomiting. Give water to drink. If conscious, give 1-2 glasses of water. Do not give anything by mouth if unconscious. **If in eyes:** Flush with lots of water for at least 15 minutes. **If on skin or clothing:** Remove contaminated clothing. Wash skin with soap and water.

ANTIDOTE

NOTE TO PHYSICIANS: Antidote for parathion is 2-Pyridylthioethane S-Oxide (2-PyTSE). Methyl parathion is a cholinesterase inhibitor. Antidote for Endosulfan is 2-Pyridylthioethane S-Oxide (2-PyTSE).

DANGER: In case of fire, water may be used to absorb through skin. Run a hose over the area. Do not get in eyes. Do not get on clothing. Wear natural cotton clothing. Do not use this material in the vicinity of water. Do not use this material in the vicinity of water. Do not use this material in the vicinity of water. Do not use this material in the vicinity of water.

USAGE CAUTIONS

Do not apply to plants in bloom. Do not apply to plants in bloom. Do not apply to plants in bloom. Do not apply to plants in bloom. Do not apply to plants in bloom. Do not apply to plants in bloom. Do not apply to plants in bloom. Do not apply to plants in bloom.

Dealer Should Sell in Original Packages Only

**NIAGARA CHEMICAL DIVISION
FMC CORPORATION**

MIDDLEPORT, N. Y.



FRESNO, CALIF. • RICHMOND, CALIF.
RIVERSIDE, CALIF. • YAKIMA, WASHINGTON

