

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and aquatic life. Do not apply to water bodies or to areas adjacent to water bodies. Do not apply to areas where it may be carried by wind or water to water bodies.

This product is highly toxic to birds and mammals. Do not apply to areas where it may be carried by wind or water to areas where birds and mammals are present.

DIRECTIONS FOR USE

This product is to be used as a foliar spray on the following crops:

Apply when insects first appear and repeat applications as necessary to maintain control. Do not apply this product in undrained form. Use the recommended rates. Use sufficient water to thoroughly cover the crop. By ground application, use 10 to 20 gallons when applying 2.0% material by airraft in a 100-foot recommended application with sufficient water to provide a minimum of 3 gallons of finished spray per acre. Observe days interval between last application and harvest indicated by number in parenthesis following the crop.

Artichoke (7) Aphids Use 1 to 2 quarts per acre. Artichoke Plant Moth Use 1 quart per acre.

Beans, Succulent and Dry (except Lima Beans) (15) Black Bean Aphid, Cucumber Beetle, Flea Beetles, Bean Leafminer, Green Stink Bug, Mexican Bean Beetle Use 1 to 2 quarts per acre. Armyworms, Leafhoppers Use 2 quarts per acre. Do not exceed 3 applications per season. Do not feed treated freshings to livestock or a low grazing treated fields.

Broccoli (7), Brussels Sprouts (14), Cabbage (21), Cauliflower (14) Aphids, Cabbage Looper, Imported Cabbageworm, Diamondback Moth Larvae, Cross-striped Cabbageworm, Flea Beetles, Stink Bugs, Harlequin Bug Use 1 to 2 quarts per acre.

Carrots (15) Flea Beetle, Green Peach Aphid Use 1 to 2 quarts per acre. Armyworm, Leafhoppers Use 2 quarts per acre. Do not exceed 2 applications per season. Do not use tops for food or feed.

Celery (Whole Plant) (15) Aphids including Green Peach Aphid, Cabbage Looper, Flea Beetles Use 1 to 2 quarts per acre. Armyworms, Leafhoppers Use 2 quarts per acre. Do not exceed 3 applications per season.

Collards (21) Cabbage Looper, Diamondback Moth Larvae, Harlequin Bug, Imported Cabbageworm Use 1 to 2 quarts per acre. Do not exceed one application per season.

Kale (21) Cabbage Flea Beetle, Imported Cabbageworm, Harlequin Bug Use 1 to 2 quarts per acre. Do not exceed one application per season.

Lettuce (21) Cabbage Looper, Aphids including Green Peach Aphid, Diamondback Moth Larvae, Imported Cabbageworm, Armyworms, Flea Beetles, Leafhoppers Use 1 to 2 quarts per acre. On head lettuce do not apply more than three times after thinning. Remove heads 30 days after harvest. Do not feed crop waste to livestock. Do not use for more than 2 applications per season.

Mustard Greens (21), Turnip Greens (21) Aphids, Cabbage Looper, Imported Cabbageworm Use 1 to 2 quarts per acre. Do not exceed one application per season.

Peas, Succulent (Northwest Only) (15) Flea Aphid, Flea Weevils Use 2 quarts per acre. Do not exceed 2 applications per season. Do not use peas when plants to be harvested by hand. Do not feed treated crop to livestock or allow to graze in treated fields.

Peppers (15) Flea Beetle, Green Peach Aphid, Pepper Maggot, Hornworm Use 1 to 2 quarts per acre. Armyworm, Leafhoppers Use 2 quarts per acre.

Spinach (21) Flea Beetle, Green Peach Aphid, Turnip Mine, Armyworm, Flea Beetles, Leafhoppers Use 1 to 2 quarts per acre. Do not exceed one application per season.

Tomatoes (10) Aphids including Green Peach Aphid, Flea Beetle, Flea Weevils Use 1 to 2 quarts per acre. Armyworm, Leafhoppers Use 2 quarts per acre. Do not use tomatoes for food or feed. Do not use for more than 2 applications per season. Do not use for more than 2 applications per season.

ACCEPTED
OCT 04 1973
U.S. DEPARTMENT OF AGRICULTURE
EPA Reg. No. 1609

STORAGE AND DISPOSAL

This product is a flammable liquid. Store in a cool, dry place. Do not store in areas where it may be exposed to fire or heat. Do not store in areas where it may be exposed to sunlight. Do not store in areas where it may be exposed to moisture.

Dealers Should Sell in Original Packages On Terms of Sale or Use

Warranty: This product is warranted to be free from defects in material and workmanship. It is not warranted to be effective against all insects and diseases.

Directions and Recommendations

Use of Product: This product is to be used as a foliar spray on the following crops: Artichoke, Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Carrots, Celery, Collards, Kale, Lettuce, Mustard Greens, Turnip Greens, Peas, Peppers, Spinach, Tomatoes.

Damages

This product is highly toxic to fish and aquatic life. Do not apply to water bodies or to areas adjacent to water bodies. Do not apply to areas where it may be carried by wind or water to water bodies.