

Code: 30823

Net Weight
50 Pounds

CRYOLITE 40 PARATHION 2 DUST

DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. May be applied by air or ground. Observe days interval between last application and harvest indicated by number () following the crop.

BEANS: Bean Leaf Beetle — Use 12½ pounds per acre. Aphids, Whitefly, Leafhopper, Mexican Bean Beetle, Bean Leafroller — Use 25 pounds per acre. Do not apply within 7 days of harvest. Corn Earworm, Cucumber Beetle — Use 37½ pounds per acre. Do not apply within 15 days of harvest. Remove excess residues by washing, brushing or other effective means.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, COLLARDS, KALE, TURNIPS, RADISH (15), and MUSTARD GREENS: Aphids, Cabbage Looper, Armyworms, Climbing Cutworms, Diamondback Moth Larvae, Imported Cabbageworm, and Flea Beetles — Use 25 pounds per acre. Vegetable Weevil — Use 20 pounds per acre. On Broccoli, Brussels Sprouts and Cabbage do not apply after the edible parts or the heads start to form. On Collards, Mustard and Kale do not apply after seedling stage. On Turnips if tops are fed do not apply after seedling stage. Remove excess residues by washing, brushing or other effective means.

CAULIFLOWER, KOHLRABI: Do not apply after seedling stage. Remove excess residues by washing, brushing or other effective means. Aphids, Climbing Cutworms — Use 25 pounds per acre. Armyworms, Diamondback Moth Larvae, Imported Cabbageworms, Vegetable Weevil — Use 20 pounds per acre. Cauliflower only: Cabbage Looper, Flea Beetles — Use 25 pounds per acre.

CUCUMBERS (15), SQUASH (15): Aphids, Climbing Cutworms, Cucumber Beetle, Squash Bug, Leafminer, Thrips and Flea Beetle — Use 25 pounds per acre. Remove excess residues by washing, brushing or other effective means. Do not apply before plants start to vine or when they are wet.

LETTUCE: Aphids, Leafhoppers, Leafminers, Cabbage Looper, Armyworm and Imported Cabbageworm — Use 25 pounds per acre. Do not apply to Bibb and Leaf Lettuce after the seedling stage. Do not apply to Head Lettuce after the head begins to form.

MELONS (7): Aphids, Leafhopper, Leafminer, Thrips, and Cucumber Beetle — Use 25 pounds per acre. Remove excess residues by washing, brushing or other effective means.


PEPPERS (15): Aphids — Use 15 pounds per acre. Remove excess residues by washing, brushing or other effective means.

INSECTICIDE

EPA REGISTRATION NO. 279-2627

ACTIVE INGREDIENTS:	BY WT.
*Sodium Fluoroaluminate	40.00%
Parathion; O,O-Diethyl O-p-nitrophenyl thiophosphate	2.00%
INERT INGREDIENTS:	58.00%
	100.00%

*Fluorine expressed as element 2.7%



DANGER - POISON



KEEP OUT OF REACH OF CHILDREN

FIRST AID

Call a Physician Immediately! If swallowed: Give a tablespoon of salt in a glass of warm water. Repeat until vomiting occurs and vomitus is clear. Have victim lie down and keep quiet. Never induce vomiting in an unconscious person.

If in Eyes: Flush eyes for 15 minutes with plenty of water.

If on Skin or Clothing: Remove contaminated clothing and shoes immediately. Bathe with soap and water.

NOTE TO PHYSICIAN: Parathion is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidotal. Give 4 mg. intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If cyanotic give artificial respiration. 2-PAM is also antidotal and may be used in conjunction with atropine.

DANGER: Poisonous if Swallowed, Inhaled or Absorbed Through Skin! Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing, and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Parathion protection. Keep all unprotected persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying at least 18 inches deep in non-crop land away from water supplies. Empty bags should be buried with waste. Do not burn.

For Emergency Assistance Call
716-735-3765