

Code: 31226

50 Pounds Net Weight

KOLODUST[®] 50 DIAZINON[®] 3 DUST

DIRECTIONS FOR USE

Apply this dust to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants. Apply the dust to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

BEETS (14) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

BROCCOLI (5), CABBAGE (7), CAULIFLOWER (5) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

CELERY (10) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

COLLARDS (10), KALE (10), PARSLEY (12), SWISS CHARD (12), TURNIP TOPS (10) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

CORN (10) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

FIGS (5) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

GRAPES (10) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

HOPS (14) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

LETTUCE (10) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

ONIONS (10) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

POLE LIMA AND SNAP BEANS (7) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

POTATOES (35) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

TOMATOES (3) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

SPINACH (10) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

STRAWBERRIES (5) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

WALNUTS (5) Apply to the foliage of the plants in the following manner: Use a backpack sprayer or a hand-held dust applicator to apply the dust to the foliage of the plants.

FUNGICIDE-INSECTICIDE

For use on the following crops: (1) Beans, (2) Corn, (3) Cucumber, (4) Eggplant, (5) Melon, (6) Pepper, (7) Pumpkin, (8) Squash, (9) Tomato, (10) Watermelon.

For use on the following crops:	For use on the following crops:
(1) Beans	(1) Beans
(2) Corn	(2) Corn
(3) Cucumber	(3) Cucumber
(4) Eggplant	(4) Eggplant
(5) Melon	(5) Melon
(6) Pepper	(6) Pepper
(7) Pumpkin	(7) Pumpkin
(8) Squash	(8) Squash
(9) Tomato	(9) Tomato
(10) Watermelon	(10) Watermelon

CAUTION

KEEP OUT OF REACH OF CHILDREN

Handle with care. Avoid contact with skin and eyes. Wash hands thoroughly after use. Avoid breathing dust. Store in a cool, dry place.

This product contains Diazinon, which is a highly toxic chemical. It is a powerful insecticide and fungicide. It is used to control insects and fungi on a variety of crops. It is a highly effective product, but it must be used with care. It is a highly toxic chemical, and it can be harmful to humans and animals if it is not used properly. It is a highly effective product, but it must be used with care.

USAGE CAUTIONS

Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label.

Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label.


Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label.

Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label.

Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label. Do not use this product on crops that are not listed on the label.

For Emergency Assistance Call
716-735-3765

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product, buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user, for which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to make any statement or warranty to the contrary.

Manufactured by
FMC CORPORATION
AGRI-INDUSTRIAL DIVISION
Philadelphia, Pa. 19101

FMC
Chemicals

Code: 31226

Net Weight

Kolodust® 50 Diazinon 3 Dust
Fungicide-Insecticide

E.P.A. REG. NO. 279-2633 AA

EPA Est., 279-

ACTIVE INGREDIENTS:	By Wt.
0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	3.00%
Sulphur	42.00%
INERT INGREDIENTS:	55.00%
	100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near food and feed products.

For Emergency Assistance Call 1-800-335-3764

See other panels for additional safety information.

Usage Precautions

This product is toxic to fish and wildlife. Keep out of lakes, ponds, or streams. Birds feeding on treated areas may be killed. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Applications of materials containing sulfur should not be made prior to or during periods of excessively high temperatures.

Do not allow this material to drift onto sulfur sensitive crops.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until dusts have settled.

Protective clothing means at least a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under "Written Warnings". Written oral warnings are given. Warnings shall be given in language clearly understood by workers. Oral warnings must be given where there is reason to believe that written warnings could not be understood by workers. Written warnings must include the following information:

CAUTION

Area treated with Kolodust® 50 Diazinon 3 Dust (in insert date). Do not enter without appropriate protective clothing until sprays have dried (insert State Department of Agriculture's re-entry interval if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

Storage and Disposal

Pesticide Storage

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animal and unprotected persons. Confine spill. Call FMC collect (766) 376-0155.

To contain spill, if liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in leaking container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

This product contains Benlate® Sulfur. Benlate® Sulfur is made by the absorption of liquid sulfur into Benlate in the proportion of thirty parts sulfur and seventy parts Benlate. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, susceptibility in spray tank and safety to host plants.

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

11/8/81
FARMER

Apply when insects or disease first appear and repeat as necessary to maintain control. For control beyond the use limitations of this product use a suitable material. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

Beets (14): Aphids, Dipterous Leafminer—Use 16 pounds per acre.

Broccoli (5), Cabbage (7), Cauliflower (5): Aphids, Diamondback Moth Larvae, Imported Cabbageworm, Powdery Mildew—Use 33-50 pounds per acre.

Celery (10): Aphids, Flea Beetles—Use 16 pounds per acre. Plants should be marked trimmed, tops removed, before being used or shipped. Tops should not be used for feed or food.

Collards (10), Kale (10), Parsley (12), Swiss Chard (12), Turnip Tops (10): Aphids, Imported Cabbageworm, Diamondback Moth Larvae—Use 40-50 pounds per acre.

Corn, Corn Earworm: Use 50-60 pounds per acre. Make first application when silks first appear. Make 2 or 3 repeat applications at 2 to 3 day intervals. Corn may be picked immediately following last application. Do not feed forage to livestock within 10 days following last application.

Figs (5): Vinegar Flies, Dried Fruit Beetles—Use 50-80 pounds per acre.

Grapes (1): Grape Leafhoppers, Powdery Mildew—Use 33 pounds per acre. Ground application preferred.

Hops (14): Beet Armyworm, Corn Earworm, Climbing Cutworm, Aphids, Cucumber Beetle—Use 65 pounds per acre.

Lettuce (10): Aphids, Dipterous Leafminer—Use 16 pounds per acre.

Onions (10): Thrips, Pea Leafminer (Pacific Northwest)—Use 40 pounds per acre.

Pole, Lima and Snap Beans (7): Powdery Mildew, Rust, Aphids, Leafhoppers, Cucumber Beetle—Use 40-45 pounds per acre. Sulfur may injure certain varieties. Check sulfur tolerance prior to use.

Potatoes (35): Aphids, Colorado Potato Beetle, Flea Beetles—Use 35-65 pounds per acre.

Tomatoes (3): Dipterous Leafminer, Vinegar Fly (Drosophila), Aphids, Russet Mite—Use 40-50 pounds per acre.

Spinach (10): Aphids, Spinach Leafminer—Use 16 pounds per acre.

Strawberries (5): Aphids, Powdery Mildew—Use 33-40 pounds per acre. Do not apply this product to Strawberries while fruit is present.

Walnuts: Walnut Acanth—Use 50-80 pounds per acre. Do not apply after husk split.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon facts believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established sale practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.