CODE
30063

Niagara®

EPN 4 EC

INSECTICIDE

NET CONTENTS
5 Gals.



DANGER-POISON

KEEP OUT OF REACH OF CHILDREN

USDA REG. NO. 279-2630

ACTIVE INGREDIENTS:

*O-Ethyl O-(p-nitrophenyl) phenylphosphonothioate	44%
Xylene Range and Aromatic Petroleum Solvent	44%
INERT INGREDIENTS	12%
	100%

*EPN

CONTAINS 4 LBS. EPN PER GALLON

DANGER: Poison by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Prolonged exposure may cause cholinesterase inhibition. Causes severe eye irritation.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash separately before reuse. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

DO NOT USE IN ENCLOSED ROOMS

DO NOT STORE BELOW 40°F

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

DIRECTIONS FOR USE

FOR THE CONTROL OF EUROPEAN CORN BORER AND CORN ROOTWORM (ADULTS) INFESTING CORN

Apply at the rate of 1 2/3 to 1 pint per acre in sufficient water to cover plant surfaces, including the leaf whorls. This material can be used by ground or aerial applicators. When applying this material by aircraft, mix the the recommended amounts with sufficient water to provide a minimum of 1 gallon of finished spray per acre. OBSERVE DAYS INTERVAL BETWEEN LAST APPLICATION AND HARVEST OR FEEDING TO LIVESTOCK AS INDICATED BY NUMBER IN ( ) FOLLOWING CROP

SWEET CORN (14): European Corn Borer - Make first application when 25% of the plants show evidence of leaf feeding and continue at 5 day intervals as long as leaf feeding is evident. From one to three treatments may be required. For control of second brood borers, apply when eggs begin to hatch. Corn Rootworm (adults) - To protect the silks from damage apply when the insects appear in sufficient number to the ripe silks and prevent pollination.

FIELD CORN (14): European Corn Borer - Apply 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is present. For control of second brood borers, apply when eggs begin to hatch. Corn Rootworm (adults) - To protect the silks from damage apply when the insects appear in sufficient number to the ripe silks and prevent pollination.

POPCORN GROWN FOR SEED (14): Apply when 25% of the plants show evidence of leaf feeding. Repeat in 7 days if feeding continues. Treat for second brood at pollen shedding time and while the plants are still in the green silk stage.

FIRST AID

If In Eyes: Flush eye for 15 minutes with plenty of water. Call a physician immediately.

If On Skin or Clothing: Remove contaminated clothing and shoes immediately. Bathe with soap and water. Call a physician immediately.

If Swallowed: Give tablespoon of table salt in a glass of warm water. Repeat until vomiting occurs and vomitus is clear. Never induce vomiting in an unconscious person. Call a physician immediately.

ANTIDOTE

Note to Physician: Active ingredient is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidotal. Give 4 mg. intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If vomiting give artificial respiration. For additional assistance call (716) 735-3761.

This product is toxic to fish and wildlife. Do not contaminate water by cleaning equipment or disposal. Do not use in Bird feeding or treated areas (not to be killed). Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging in the area. Agricultural Extension Service.

Rinse equipment and containers and DISPOSE OF WASTES by burying at least 18 inches deep in non-crop lands away from water supplies. Do not reuse container or dispose of in any manner.

DO NOT APPLY THIS MATERIAL TO DIETARY OR NURSERY CROPS OR STOCK FOR DAIRY USE. DO NOT APPLY DIRECTLY TO MAN OR ANIMAL OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS. AVOID CONTACT WITH SKIN, EYES OR CLOTHING. WEAR PROTECTIVE CLOTHING WHILE HANDLING OR APPLYING THIS MATERIAL.

Dealers Should Sell In Original Packages Only.

TERMS OF SALE OR USE: Buyer assumes the risks of use or handling of this product not in accordance with the Directions for Use.

WARRANTY: Niagara warrants that the product is of the kind described in the accompanying statement on the label and that it is suitable for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to the purchase price paid and shall not include any other special or consequential damages. Notice of any claim must be given to Niagara within 10 days after such claim shall have arisen or be barred. Failure to give such notice shall constitute a waiver by buyer of such claim.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of various factors affecting the use of this product, such as weather conditions, method or manner of use, presence of other materials or other factors in connection with which Niagara has no control, buyer and user assume all responsibility for any loss or injury resulting from the handling or use of the product not attributable to fault on the part of Niagara. No person is authorized to vary or waive any statement contained herein.

Printed in U.S.A.

030