

Code 30063

Net Contents

Gals.

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification

BEST DOCUMENT AVAILABLE

EPN 4 EC

Insecticide

For Agricultural or Commercial Use Only.

EPA Reg. No. 279-2650

EPA Est.

Active Ingredients

O-Ethyl O-p-nitrophenyl phenylphosphonothioate	45.5%
Xylene base aromatic petroleum solvent	40.9%
Inert Ingredients	13.6%

EPN 100.0%
Contains 4 lbs EPN Per Gallon

First Aid

If Swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If In Eye, flush eye for 15 minutes with plenty of water. Call a physician immediately.

If On Skin or Clothing, remove contaminated clothing and shoes immediately. Bathe with soap and water. Call a physician immediately.

Antidote

Note to Physicians: Active ingredient is a cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine antidotal. Give 4 mg intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If cyanotic give artificial respiration.

For Emergency Assistance Call 716-735-3765.

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Danger

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Repeated exposure may, without symptoms, be increasingly hazardous.

Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before reuse. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and warm water; for eyes, flush with water for 15 minutes and get medical attention.

Do not get in eyes, on skin, or on clothing. Do not breathe sprays. Protective Clothing Required. Wear clean protective clothing, goggles and respirator approved by NIOSH or the American National Standards Institute when applying or handling, or when reentering fields within a 24-hour period. The following protective clothing must be worn: lightweight, lined or natural rubber gloves, at least mid forearm length; well-sealed waterproof hat or waterproof hood; a protective suit or coveralls of a nonpermeable, non-cloth material covering the body from ankles to wrists; lightweight, unlined, natural rubber boots at least midcalf in length. Full-face respirators are recommended; half-face respirators and goggles are required. Aerial applicators in positive pressure suits and helmet applicators in comparable ground equipment with appropriate filters at all air intakes need not comply with these protective clothing requirements.

Environmental Hazards

This product is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Birds feeding in treated areas may be killed. Apply the product only as specified on this label.

Do not apply this product to areas where bees are visiting the treated area.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, legal liability or other undesirable results may occur.


Physical or Chemical Hazards

None known.

DIRECTIONS FOR USE

For use on... (text partially obscured)

KEEP OUT OF REACH OF CHILDREN



PELIGRO

PRECAUCION AL USUARIO: Se listados en etiquetas, no use este producto...

See Other Panels for First Aid and Precautionary Information.



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

Market Order PA 99902 Release number WG 1111

Do not apply this product in a manner as to directly or through drift injure or damage crops, animals, or humans.

Do not apply this product in a manner as to directly or through drift injure or damage crops, animals, or humans.

Written oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing. Periods of time fields must be vacated and appropriate safety measures taken to avoid exposure. An example of such a warning is given below. Written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

DANGER

Area treated with EPA # EC on (insert date). Do not enter without appropriate protective clothing. (R-24) (EPA # 5 (insert State Department of Agriculture's re-entry interval) (more restrictive).

In case of accidental exposure to pesticide spray or dust, wash immediately with copious amounts of water. Remove contaminated clothing and wash before reuse. If necessary, flush with plenty of water. If needed, go to an area where the pesticide has not been applied. Get medical attention if needed.

Storage and Disposal

Pesticide Storage

Do not store below 50°F (10°C).

Do not use or store near heat, open flame or hot surfaces.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place, away from direct sunlight. Carefully open containers. Avoid contact with skin and eyes. Do not put concentrate or dilute material in food or drink containers. Do not contaminate other pesticides, fertilizers, water, or other agricultural materials.

For more information, contact FMC, 10000 North Central Expressway, Dallas, Texas 75243, or 1-800-375-3755.

The contents of this container are not to be used for any other purpose. May be sold or disposed of only in accordance with the instructions on the label. Do not use for any other purpose.

Pesticide Disposal

Pesticide wastes are acutely toxic. Do not discharge into the environment. Do not use for any other purpose. Do not use for any other purpose. Do not use for any other purpose. Do not use for any other purpose.

Container Disposal

Do not use or equipment. Then offer for recycling or reuse. Do not use for any other purpose. Do not use for any other purpose. Do not use for any other purpose. Do not use for any other purpose.

Do not apply this product when weather conditions favor drift from treated areas. Do not apply to corn during the pollen shed period if bees are present in the area.

Apply at the rate of 1.0 to 1.5 pints per acre in sufficient water to cover plant surfaces, including the leaf whorls. This material can be used by ground or air application. When applying this material by aircraft, use the maximum label amounts with sufficient water to provide a minimum of 1 gallon of finished spray per acre. Observe days interval between last application and harvest or feeding to livestock as indicated by number in the following crop.

Apply at the rate of 1.0 to 1.5 pints per acre in sufficient water to cover plant surfaces, including the leaf whorls. This material can be used by ground or air application. When applying this material by aircraft, use the maximum label amounts with sufficient water to provide a minimum of 1 gallon of finished spray per acre. Observe days interval between last application and harvest or feeding to livestock as indicated by number in the following crop.

Sweet Corn (14): European Corn Borer—Make first application when 25% of the plants show evidence of leaf feeding and continue at 5 day intervals as long as leaf feeding is evident. From one to three treatments may be required. For control of second brood borers, apply when eggs begin to hatch. Corn Rootworm (adult)—To protect the silks from damage apply when the insects appear in sufficient number to damage silks and prevent pollination.

Field Corn (14): European Corn Borer—Apply 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is present. For control of second brood borers, apply when eggs begin to hatch. Corn Rootworm (adult)—To protect the silks from damage apply when the insects appear in sufficient number to damage silks and prevent pollination.

Corn Grown for Seed (14): Apply when 25% of the plants show first brood whorl feeding. Repeat in 7 days if feeding continues. Treat for second brood at pollen shedding time and while the plants are still in the green silk stage.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions.

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be relative. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

FMC

BEST DOCUMENT AVAILABLE

Ms. Eunice B. Curle
FMC Corporation
2000 Market Street
Philadelphia, PA 19103

Dear Ms. Curle:

Subject: EPN 42C
EPA Registration No. 279-2656
response to EPN L.A. Notice of August 31, 1963
Your Letter Dated October 25, 1963

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

However, before you release the product for shipment, make the following change.

1. Submit your final printed labeling as follows:

for aerial application, flagging must be by fully automated mechanical means or by humans working in totally enclosed vehicles.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (45-767)

IRB/RD:ELLENBERGER:DCR-21436:WABG-1264C:mk:kaven:479-2013:11/18/83

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

EPA Form 1320 (4-81) OFFICIAL FILE COPY