

Code: 30117

MANUFACTURED:

SULPHUR 50 DIBROM[®] 4 SEVIN[®]* 5 DUST

GU

DIRECTIONS FOR USE

Apply when pests or disease first appear and repeat as necessary to maintain control. For control beyond the use limitations of this product use a suitable material. May be applied by air or ground. Observe days interval between last application and harvest indicated by number in () following the crop.

BROCCOLI (3), BRUSSELS SPROUTS (3), CABBAGE (3), CAULIFLOWER (3): Flea Beetle, Harlequin Bug — Use 10 to 20 pounds per acre. Armyworm, Imported Cabbageworm, Diamondback Moth Larvae, Corn Earworm — Use 20 to 40 pounds per acre. Cabbage Looper, Aphids — Use 40 pounds per acre. For small Cabbage Looper control, after edible parts start to form apply at 5 to 7 day intervals.

COLLARDS (14), KALE (14), TURNIP GREENS (14): Flea Beetle, Harlequin Bug, Leafhoppers — Use 10 to 20 pounds per acre. Armyworm, Corn Earworm, Tarnished Plant Bug, Stink Bugs, Imported Cabbageworm, Diamondback Moth Larvae — Use 20 to 40 pounds per acre. Cabbage Looper, Aphids — Use 40 pounds per acre. For small Cabbage Looper control, after edible parts start to form, apply at 5 to 7 day intervals.

COTTON (4): Early Season: Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Striped Blister Beetle — Use 10 to 20 pounds per acre. Grasshopper — Use 10 to 30 pounds per acre. Mid and Late Season: Boll Weevil, Bollworm, Fall Armyworm, Cotton Leafperforator, Leafrollers, Leafhoppers, Tarnished Plant Bug, Lygus Bug, Aphids (suppression) — Use 20 to 35 pounds per acre. Pink Bollworm — Use 30 to 35 pounds per acre. Armyworm, Cabbage Looper — Use 35 pounds per acre.

GRAPES (4): Powdery Mildew, Grape Leafhopper, Grape Leafroller, Grape Leaf-skeletonizer, Thrips — Use 20 to 40 pounds per acre. Concord and other Labrusca type grapes may be injured by sulfur. Ground application preferred.

LETTUCE (Head-3, Leaf-14), SPINACH (14), SWISS CHARD (14): Flea Beetle, Harlequin Bugs, Leafhoppers — Use 10 to 20 pounds per acre. Armyworms, Corn Earworm, Tarnished Plant Bug, Stink Bugs, Imported Cabbageworm — Use 20 to 40 pounds per acre. Aphids, Cabbage Looper — Use 40 pounds per acre.

PEACHES (4) (Western States): Powdery Mildew, Brown Rot, Aphids, Fruit Fly, Red Spider Mites, Lygus Bug, Thrips, Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth — Use 60 to 80 pounds per acre. For Codling Moth control apply at petal fall and every 10 to 14 days thereafter.

PEPPERS (1): Colorado Potato Beetle, Flea Beetle, Leafhoppers — Use 10 to 20 pounds per acre. Fall Armyworm, Tomato Hornworm, Tomato Fruitworm, Tarnished Plant Bug, Stinkbug — Use 20 to 25 pounds per acre. Aphids, Leafminers — Use 25 pounds per acre.

STRAWBERRIES (4): Meadow Spittlebug, Strawberry Leafroller, Powdery Mildew, Strawberry Weevil — Use 20 to 40 pounds per acre. Red Spider Mites, Aphids — Use 40 pounds per acre.

TOMATOES (1): Colorado Potato Beetle, Flea Beetles, Leafhoppers — Use 10 to 20 pounds per acre. Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug, Stink Bug, Tomato Russet Mite, Fruit Fly, Twospotted Spider Mite, Aphid, Leafminer — Use 25 pounds per acre.

FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279 2657 ZA

ACTIVE INGREDIENTS	BY WT.
*Carbaryl, 1-naphthyl N-methylcarbamate	5.00%
**Naled, 1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	4.00%
Sulphur	50.00%
INERT INGREDIENTS	41.00%
	100.00%

*Trademark Union Carbide Corporation for the active ingredient carbaryl.

**DIBROM is a registered trademark of Chevron Chemical Company. U.S. Patent No. 2,971,882.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. May cause irritation of eyes, nose and throat. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL AND

This product is toxic to fish, birds and other streams. Birds feeding on treated areas may be affected. Clean equipment, or disposal of waste materials favor drift from the areas treated. Do not occur. Apply this product only as specified.

This product is highly toxic to bees exposed to it. Protective information may be obtained from Extension Service.

DISPOSE OF WASTES by burying in non-hazardous waste. Empty bags should be buried with wastes.

Some phytotoxicity may occur on tender foliage of several days duration following application. Foliage injury to certain crops.

Applications of materials containing sulfur should be avoided during periods of excessively high temperatures. Do not treat.


STORE IN COOL, DRY PLACE.

May corrode metal dust equipment; clean equipment.

When growing crops for processing, consult

Consult with your State Agricultural Experiment Station Extension Service for additional information and directions.

For Emergency Assistance Call: 716-735-3765



Net Weight
50 Pounds

GUARANTEE EXPIRES:

PHUR 50 DIBROM[®] 4 SEVIN[®]* 5 DUST

FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279-2657 ZA

ACTIVE INGREDIENTS:		BY WT.
*Carbaryl; 1-naphthyl N-methylcarbamate		5.00%
**Naled; 1,2-dibromo-2,2-dichloroethyl dimethyl phosphate		4.00%
Sulphur		50.00%
INERT INGREDIENTS:		41.00%
		100.00%

*Trademark Union Carbide Corporation for the active ingredient carbaryl.
**DIBROM is a registered trademark of Chevron Chemical Company.
U.S. Patent No. 2,971,882.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. May cause irritation of eyes, nose and throat. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application. Sulfur may cause severe fruit and foliage injury to certain crops.

Applications of materials containing sulfur should not be made prior to or during periods of excessively high temperatures. Do not use within 2 weeks of an oil spray treatment.


STORE IN COOL, DRY PLACE.

May corrode metal dust equipment; clean equipment after use.

When growing crops for processing, consult processor before applying sulfur.

Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with these label cautions and directions.

For Emergency Assistance Call: 716-735-3765



EPA EST., 279-CA-1 Printed in U.S.A.

Code: 30117

MANUFACTURED:

SULPHUR 50 DIBROM

DIRECTIONS FOR USE

Apply when pests or disease first appear and repeat as necessary to maintain control. For control beyond the use limitations of this product use a suitable material. May be applied by air or ground. Observe days interval between last application and harvest indicated by number in following the crop.

BROCCOLI (3), BRUSSELS SPROUTS (3), CABBAGE (3), CAULIFLOWER (3): Flea Beetle, Harlequin Bug. Use 10 to 20 pounds per acre. Armyworm, Imported Cabbageworm, Diamondback Moth Larvae, Corn Earworm. Use 20 to 40 pounds per acre. Cabbage Looper, Aphids. Use 40 pounds per acre. For small Cabbage Looper control, after edible parts start to form, apply at 5 to 7 day intervals.

COLLARDS (14), KALE (14), TURNIP GREENS (14): Flea Beetle, Harlequin Bug, Leafhoppers. Use 10 to 20 pounds per acre. Armyworm, Corn Earworm, Tarnished Plant Bug, Stink Bug, Imported Cabbageworm, Diamondback Moth Larvae. Use 20 to 40 pounds per acre. Cabbage Looper, Aphids. Use 40 pounds per acre. For small Cabbage Looper control, after edible parts start to form, apply at 5 to 7 day intervals.

COTTON (4) Early Season: Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Striped Bister Beetle. Use 10 to 20 pounds per acre. Grasshopper. Use 10 to 30 pounds per acre. **Mid and Late Season:** Boll Weevil, Bollworm, Fall Armyworm, Cotton Leafperforator, Leafrollers, Leafhoppers, Tarnished Plant Bug, Lygus Bug, Aphids (suppression). Use 20 to 35 pounds per acre. Pink Bollworm. Use 30 to 35 pounds per acre. Armyworm, Cabbage Looper. Use 35 pounds per acre.

GRAPES (4): Powdery Mildew, Grape Leafhopper, Grape Leafroller, Grape Leaf skeletonizer, Thrips. Use 20 to 40 pounds per acre. Concord and other Labrusca type grapes may be injured by sulfur. Ground application preferred.

LETTUCE (Head-3, Leaf-14), SPINACH (14), SWISS CHARD (14): Flea Beetle, Harlequin Bugs, Leafhoppers. Use 10 to 20 pounds per acre. Armyworms, Corn Earworm, Tarnished Plant Bug, Stink Bugs, Imported Cabbageworm. Use 20 to 40 pounds per acre. Aphids, Cabbage Looper. Use 40 pounds per acre.

PEACHES (4) (Western States): Powdery Mildew, Brown Rot, Aphids, Fruit Fly, Red Spider Mites, Lygus Bug, Thrips, Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth. Use 60 to 80 pounds per acre. For Codling Moth control apply at petal fall and every 10 to 14 days thereafter.

PEPPERS (1): Colorado Potato Beetle, Flea Beetle, Leafhoppers. Use 10 to 20 pounds per acre. Fall Armyworm, Tomato Hornworm, Tomato Fruitworm, Tarnished Plant Bug, Stinkbug. Use 20 to 25 pounds per acre. Aphids, Leafminers. Use 25 pounds per acre.

STRAWBERRIES (4): Meadow Spittlebug, Strawberry Leafroller, Powdery Mildew, Strawberry Weevil. Use 20 to 40 pounds per acre. Red Spider Mites, Aphids. Use 40 pounds per acre.

TOMATOES (1): Colorado Potato Beetle, Flea Beetles, Leafhoppers. Use 10 to 20 pounds per acre. Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug, Stink Bug, Tomato Russet Mite, Fruit Fly, Twospotted Spider Mite, Aphid, Leafminer. Use 25 pounds per acre.

FUNGICIDE

EPA REGISTRATION

ACTIVE INGREDIENTS:

*Carbaryl, 1-naphthyl N-methyl

**Naled; 1,2 dibromo-2,2 dichloro

Sulphur

INERT INGREDIENTS:

*Trademark Union Carbide Carbaryl.

**DIBROM is a registered trademark U.S. Patent No. 2,971,882


CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. May be harmful. Avoid contact. Wash with soap and water. Shower or bath after work. Wash and change to clean clothing daily.

AVOID CONTAMINATION OF WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a cholinesterase inhibitor. Atropine is antidotal.



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.

LABEL

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals