

Code: 30895

[279-2699]

ETHION 4 SEVIN® 10 DUST

DIRECTIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in () following the crop.

ALMONDS (30): Codling Moth, Filbert Worm, Fruitree Leafroller, Brown Mite, European Red Mite, Twospotted Spider Mite — Use 40 to 50 pounds per acre. Do not apply more than once during the fruiting period. A rate of 51 to 60 pounds per acre may also be used if no application is made after husk split.

BEANS (2): Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Erika Beetles, Leafhoppers, Corn Earworm, Stink Bug, Tarnished Plant Bug, Thrips, Twospotted Spider Mite — Use 20 pounds per acre. Do not feed treated foliage to livestock.

COTTON: Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Striped Blister Beetle, Boll Weevil, Bollworm, Fall Armyworm, Cotton Leafroller, Leafrollers, Leafhoppers, Tarnished Plant Bug, Lygus Bug, Aphids, Cotton Rootworm, Grasshoppers, Stink Bug and Saltmarsh Caterpillar — Use 25 pounds per acre. Do not apply after bolls begin to open. Cotton that has been treated should not be grazed or fed to dairy animals or other livestock. In California, workers entering treated fields within 24 hours following application should wear protective clothing.

GRAPES: Grape Leafhopper, Twospotted Spider Mite, Wallemite Mite, Grape Leafroller, Grape Leafminer — Use 20 to 30 pounds per acre. Make no more than 2 applications nor apply within 15 days of harvest, or make no more than 3 applications nor apply within 30 days of harvest. Do not apply to Salvador, Muscat or Fehrszago varieties of grapes as injury may occur. Ground application preferred.

NECTARINES (30), PEACHES (30): Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth, European Red Mite, Twospotted Spider Mite, Brown Mite — Use 40 to 60 pounds per acre. Do not apply more than twice during the fruiting period.

PLUMS, PRUNES: Peach Twig Borer, Leafhoppers, Codling Moth, Orange Tortrix, Western Tussock Moth, Brown Mite, European Red Mite, Twospotted Spider Mite — Use 40 to 60 pounds per acre. At the 40 to 43 pounds per acre rate, do not make more than 3 applications during the fruiting period nor apply within 21 days of harvest. At the rate of 44 to 60 pounds per acre, do not make more than 2 applications during the fruiting period, nor apply within 45 days of harvest.

TOMATOES (2): Colorado Potato Beetle, Flea Beetles, Leafhoppers, Fall Armyworms, Tomato Fruitworms, Tomato Hornworms, Tarnished Plant Bugs, Stink Bugs, Tropical Spider Mites, Twospotted Spider Mites, Leafminers — Use 20 pounds per acre.

INSECTICIDE — MITICIDE

EPA REG. NO. 279 2699 AA

ACTIVE INGREDIENTS	BY WT.
Ethion, 0,0,0 0-Tetraethyl S,S'-methylene diphosphorodithiate	4.00%
*Carbaryl, 1-naphthyl N-methylcarbamate	10.00%
INERT INGREDIENTS	85.00%
	<u>100.00%</u>

U.S. Patent No. 2,873,228

ENVIRONMENTAL AN

This product is toxic to fish and wildlife. Do not apply when weather conditions favor where run-off is likely to occur. Do not use for disposal of wastes. Apply this product

This product is highly toxic to bees. Protective information may be obtained from the Service.

DISPOSE OF WASTES by burying in non-combustible bags should be buried with wastes or

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. May be absorbed through skin. Avoid breathing dust. Avoid contact with eyes or skin. Wash thoroughly after handling. Wash all contaminated clothing with soap and water before re-use. Avoid storage near or contamination of feed and food-stuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is anticholinergic.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

For Emergenc
716-

Trademark of Union Carbide Corporation

279-2699

ETHION 4 SEVIN® 10 DUST



Net Weight
50 Pounds

FOR USE

Necessary to maintain control. May be applied in last application and harvest indicated by

Fruitree Leafroller, Brown Mite, European
50 pounds per acre. Do not apply more
of 51 to 60 pounds per acre may also be

Beetle, Cucumber Beetles, Flea Beetles,
Two-spotted Spider Mite
ed foliage to livestock

per, Cotton Leafworm, Striped Blister Beetle,
Leafroller, Leafrollers, Leafhoppers,
Passion, etc. Bollworm, Grasshoppers, Stink
as per acre. Do not apply after bolls begin
not be grazed or fed to dairy animals or
treated fields within 24 hours following

per Mite, Williamsite Mite, Grape Leafhopper,
per acre. Make no more than 2 applications
no more than 3 applications nor apply
Salvador, Muscat or Fehr-Szago varieties of
n preferred

per Moth, Peach Twig Borer, Periodical Cicada,
Two-spotted Spider Mite, Brown Mite —
one than twice during the fruiting period

ers, Codling Moth, Orange Tortrix, Western
Two-spotted Spider Mite — Use 40 to 60
acre rate, do not make more than 3 applica-
21 days of harvest. At the rate of 44 to 60
applications during the fruiting period, nor

beetles, Leafhoppers, Fall Armyworms, Tomato
Bugs, Stink Bugs, Tropical Spider Mites,
0 pounds per acre

INSECTICIDE — MITICIDE

EPA REG. NO. 279 2699 AA

ACTIVE INGREDIENTS	BY WT
Ethion, 0,0,0'-O'-Tetraethyl S,S'-methylene bisphosphorodithioate*	4.00%
*Carbaryl; 1-naphthyl N-methylcarbamate	10.00%
INERT INGREDIENTS	86.00%
	<u>100.00%</u>

U.S. Patent No. 2,873,228

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. May be absorbed through skin. Avoid breathing dust. Avoid contact with eyes or skin. Wash thoroughly after handling. Wash all contaminated clothing with soap and water before re-use. Avoid storage near or contamination of feed and food-stuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep Out Of Smoke.

For Emergency Assistance Call:
716-735-3765

1. Trademark of Union Carbide Corporation

279-2699

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions
WARRANTY: FMC warrants that this product, as of the time of sale by FMC (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.
DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.
EXCUSE FROM NON CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.



Manufactured in U.S.A. by
FMC CORPORATION
Agricultural Chemical Division
Millsboro, N. J. 08105

FMC
Chemicals