

AUG 15 1973

Pesticides Regulation

THIODAN®

Viagara

NET CONTENTS
5 Gallons

DIRECTIONS FOR USE

INSECTICIDE

EPA REG NO 279.2700

ACTIVE INGREDIENTS

2-Ethyl-1,3-bis(4-hydroxyphenyl)butane	24.00
2,4,6-Trinitrophenol (Picric acid)	1.80
Pyrothion	0.18
Xylene	69.02
TOTAL SOLIDITIES	5.00

U.S. Patent No. 2,590,685

1. *Prüfung der Sachverhalte* (10 Punkte)
 2. *Prüfung der Sachverhalte* (10 Punkte)
 3. *Prüfung der Sachverhalte* (10 Punkte)
 4. *Prüfung der Sachverhalte* (10 Punkte)
 5. *Prüfung der Sachverhalte* (10 Punkte)
 6. *Prüfung der Sachverhalte* (10 Punkte)
 7. *Prüfung der Sachverhalte* (10 Punkte)
 8. *Prüfung der Sachverhalte* (10 Punkte)
 9. *Prüfung der Sachverhalte* (10 Punkte)
 10. *Prüfung der Sachverhalte* (10 Punkte)

¹³ product contains 2 pounds Thiodan per gallon

WARNING

KEEP OUT OF REACH OF CHILDREN.

For Emergency Assistance Call 716-735-3765

ANTIDOTE

Notes: Physostigmine (Tropicamide) is a central nervous system stimulant and is a partial antidote. There is no specific antidote. Barbituric acid derivatives may be used in treatment.

WARNING: Hazardous if swallowed, inhaled, or absorbed through skin.
 Do not get the spray in your eyes, on skin or on clothing. In
 case of contact with eyes, flush eyes with water for 15 minutes. In
 case of contact with skin, wash with soap and water. Wash thoroughly
 after use. Do not breathe vapors or spray. Do not get in eyes, on skin, or
 on clothing. Do not use near open flame. Do not use in spray painting and
 do not use in enclosed areas. Do not use in confined spaces. Do not use in
 areas where there is a fire hazard. Do not use in areas where there is a
 risk of explosion. Do not use in areas where there is a risk of fire.
 Do not use in areas where there is a risk of explosion. Do not use in
 areas where there is a risk of fire. Do not use in areas where there is a
 risk of explosion. Do not use in areas where there is a risk of fire.

WARNING: Workers entering treated areas within 24 hours of application wear protective clothing.

DO NOT STORE AT TEMPERATURES BELOW 20° F.

DO NOT USE IN UNDILUTED FORM

[illegible]

Do not use herbicides that should not be applied when bees are foraging the area.

For spraying and containers and **DISPOSE OF WASTES** by burying at least 6 inches deep in non-crop lands away from water supplies. Do not reuse container, wash or perforate and bury with waste.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

Thosol is a registered trademark of Cansolco Hoechst Limited

FMC
Chemicals

SPINACH (21) Crown Mite, Green Peach Aphid 1 1/2 qts. Do not exceed one application

STRAWBERRIES: (4) Cyclamen Mite - 4 qts. per acre in 100 gals. of water. Do not apply at intervals of less than 35 days when fruit is present. Meadow Spittlebug, Strawberry Aphid, Tarnished Plant Bug - 2 qts. per acre in 200 gals. of water. Do not apply at intervals of less than 15 days or more than twice during any 35 day period when fruit is present.

SUGAR BEETS 30 Green Peach Aphid 1 to 2 qts. Meadow Spittlebug
1 qt. Sugar Beet Webworm 1 to 2 qts. Do not feed treated tops
to livestock

SWEET CORN - In h vegetable use only. **16 Corn Leaf Aphid** - 2 qts. **Corn Earworm** - 5 qts. Apply at 2 to 3 day intervals beginning when silks first appear and continuing until they begin to dry. Do not exceed 5 applications. Do not apply to sweet corn to be processed. Do not feed treated forage or ensilage to livestock or allow to graze in treated field.

SWEET POTATOES Sweet Potato Flea Beetle 1 qt. per acre. Apply shortly after transplanting or as soon as flea beetles appear. If necessary, make a second application 7 to 10 days later.

TOBACCO 5 Flea Beetle, Hornworm, Budworm, Green Peach Aphid, Cabbage Looper. **FIELD** 1 to 2 qts. **SEED BED** Use 1 qt. per 100 gals. of water and apply 6 gals. spray per 10 to 20 yds. **Green June Bug** Larvae. Use 1 qt. per 100 gals. of water and dump plant material 1 to 2 gals. per sq. ft. **St. J. Fire** Use 1 qt. per 100 gals. of 2 to 3 qt.

TOMATOES (Fruit & Green) Water 1 qt., Colorado 1 qt.
 Beetle Flies 1 qt. Colorado 1 qt.
 Horned Cuscutt Mite 1 qt. Colorado 1 qt.
 Stink Bugs 1/2 to 2 qts. Colorado 1 qt. Colorado 1 qt.
 per 100 gals. in 100 to 200 gals. water 1 qt. Whitefly = 1 qt.

ROOT CROPS—Dried roots of sweet potatoes (*Solanum tuberosum*) are used as a source of starch. The roots are washed, peeled, sliced, and dried. They are then ground into a fine powder.

Dealers Should Sell in Original Packages Only

TERMS OF SALE OR USE: The information on this page is provided by the advertiser and is not intended to be used for any purpose other than that for which it was intended. The advertiser is not responsible for any errors or omissions in the information provided. The advertiser is not responsible for any damages or losses resulting from the use of the information provided. The advertiser is not responsible for any damages or losses resulting from the use of the information provided.

[illegible]

EXAMINES: Basic and advanced written and oral examinations, including the patches, paper, and tape, quarterly tests, and a final exam 10 days of discovery, the final exam, and every day for the damage.

[illegible]