

CODE  
**20488**

# G-5 GRANULES

NET WEIGHT  
**50 lbs.**

## INSECTICIDE

DIRECTIONS FOR USE  
FOR USE IN LOUISIANA ONLY.

CROP: SUGAR CANE


TO CONTROL: Sugar Cane Borer

Use 20 pounds of granules per acre, applied as a broadcast application. Do not apply more than 5 times per season. Do not apply within 30 days of harvest.

May be applied by aircraft at any time during the day, provided wind conditions are relatively calm. Aircraft should be flown at a height that will insure uniform coverage. Consult your State Agricultural Extension Service or Experiment Station for specific information regarding timing of application.

Do not feed treated forage or harvest trash to livestock. Do not use on other crops used for food or forage.

Bagasse from Guthion treated sugar cane can be used for feed for cattle, sheep and goats.



\*Guthion is a Reg. TM in U. S. and Canadian Pat. Offs by Ferbenfabriken Bayer A. G., Chemagro Corporation licensee Printed in U. S. A.

E. P. A. Reg. No. 279-2778  
ACTIVE INGREDIENTS:  
\*O, O-Dimethyl S-[4-oxo-1, 2, 3-benzotriazin-3-(4-H)-ylmethyl] phosphorodithioate ----- 5.0%  
INERT INGREDIENTS: ----- 95.0%  
100.0%  
\*Guthion U. S. Patent No. 2,758,115;  
Canadian Patent No. 552,660

Use Before: (Date of expiration should not be more than 40 days from manufacture)

### WARNING

KEEP OUT OF REACH OF CHILDREN

May be poisonous if swallowed, inhaled or absorbed through the skin. Harmful if inhaled or absorbed through skin. Do not get in eyes on or skin. Do not breathe dust. Wear protective gloves when handling. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed or food.

### FIRST AID TREATMENT

Should poisoning occur, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Guthion is a cholinesterase inhibitor and treatment of poisoning should include the use of atropine.

### ATROPINE IS ANTIDOTAL

Keep on hand an emergency supply of 20 tablets, grains 1/100. In case of poisoning take 3 tablets (grs. 1/100) and call physician at once.

PHYSICIAN: Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary. **USAGE CAUTIONS**

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Dispose of wastes by burying at least 18" deep in non-crop lands away from water supplies. Do not re-use container. Smash and bury with wastes. **DO NOT BURN.**