

Code 20488

Pounds Net Weight

# G-5 Granules Insecticide

For Agricultural or Commercial Use Only

For Use in Louisiana Only

EPA Reg. No. 279-2778      EPA Est., 279-

Active Ingredient:	By Wt.
*O,O-Dimethyl S-[4-oxo-1,2,3-benzotriazin-3-(4-H)-ylmethyl] phosphorodithioate . . . .	5.0%
Inert Ingredients: . . . . .	95.0%
	100.0%

\*Guthion is a Registered Trademark in U.S. and Canadian Pat. Offs by Farbenfabriken Bayer A.G., Chemagro Corp. licensee.

Canadian Patent No. 552,660.

Use Before: (Date of expiration should not be more than 40 days from manufacture).

**KEEP OUT OF REACH OF CHILDREN**

## WARNING

### First Aid Treatment

Should poisoning occur, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

**Note to Physician:** Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

Guthion is a cholinesterase inhibitor and treatment of poisoning should include the use of atropine.

For Emergency Assistance Call 716-735-3785.

See Other Panels for Additional Precautionary Information.

# FMC

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

PA 12/80

### Atropine is Antidotal

Keep on hand an emergency supply of 20 tablets, grains 1/100. In case of poisoning take 3 tablets (grs 1/100) and call physician at once.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans (& Domestic Animals)

#### Warning

May be poisonous if swallowed, inhaled or absorbed through the skin. Harmful if inhaled or absorbed through skin. Do not get in eyes or on skin. Do not breathe dust. Wear protective gloves when handling. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

### Environmental Hazards

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in a landfill approved for pesticide containers, or bury in a safe place. Do not burn this container. Do not reuse this container. Consult Federal, State or local disposal authorities for approved alternative procedures.



**Crop: Sugar Cane****To Control: Sugar Cane Borer**

May be applied by aircraft at any time during the day, provided wind conditions are relatively calm. Aircraft should be flown at a height that will insure uniform coverage. Consult your State Agricultural Extension Service or Experiment Station for specific information regarding timing of application.

Use 20 pounds of granules per acre, applied as a broadcast application. Do not apply more than 5 times per season. Do not apply within 30 days of harvest.

Do not feed treated forage or harvest trash to livestock. Do not use on other crops used for food or forage.

Bagasse from Guthion treated sugar cane can be used for feed for cattle, sheep and goats.

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

(20488-12/80-PA)